For Research Use Only

Marcks Polyclonal antibody

Catalog Number:10004-2-lg Featured Product 14 Publications

BC046601

17118

P26645 Full Name:

GeneID (NCBI):

UNIPROT ID:

GenBank Accession Number:



Basic Information

Catalog Number: 10004-2-lg Size: 2000 μg/ml Source:

Rabbit Isotype:

myristoylated alanine rich protein kinase C substrate

Calculated MW: 39 kDa Observed MW: 88 kDa

Purification Method: Protein A purification Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500

Applications

Tested Applications: WB, IHC, ELISA **Cited Applications:** WB, IHC, IF, IP Species Specificity: mouse

Cited Species: mouse, rat, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse brain tissue. IHC: mouse pancreas tissue,

Background Information

MARCKS is a member of MARCKS family of PKC substrate and it is also a widely accepted indicator of PKC activation. It shows a ubiquitous tissue distribution. The calculated molecular weight of MARCKS is 39 kDa and its apparent molecular weights were found at 68 kDa to 90 kDa in different species. This protein is a known to regulated cytoskeleton remodeling and membrane recycling. The anti-MARCKS antibody is raised against purified MARCKS protein from mouse brain using a method similar to that previously reported for MacMARCKS

Notable Publications

Author	Pubmed ID	Journal	Application
Zhenzhen Chen	29588366	J Biol Chem	WB,IHC
Qing-Yan Zhang	35850487	Mol Cell Endocrinol	WB
Mingyue Zhong	35726797	Food Funct	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunohistochemical analysis of paraffinembedded mouse pancreas tissue slide using 10004-2-Ig (Marcks antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse pancreas tissue slide using 10004-2-Ig (Marcks antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse brain tissue were subjected to SDS PAGE followed by western blot with 10004-2-Ig (Marcks antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.