

For Research Use Only

cIAP1 Polyclonal antibody

Catalog Number: 10022-1-AP

Featured Product

21 Publications



Basic Information

Catalog Number:

10022-1-AP

Size:

800 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG15203

GenBank Accession Number:

BC016174

GeneID (NCBI):

329

UNIPROT ID:

Q13490

Full Name:

baculoviral IAP repeat-containing 2

Calculated MW:

618 aa, 70 kDa

Observed MW:

70 kDa, 55-60 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IHC 1:50-1:500

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse

Cited Species:

human, mouse, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse skeletal muscle tissue, Jurkat cells, mouse brain tissue, mouse liver tissue, mouse testis tissue

IHC : human testis tissue, mouse pancreas tissue, mouse testis tissue, human pancreas tissue

IF/ICC : HepG2 cells,

Background Information

BIRC2 (also known as cIAP1) is a member of the inhibitor of apoptosis protein (IAP) family. The inhibitor of apoptosis (IAP) proteins are a family of anti-apoptotic regulators found in viruses and metazoans. BIRC2 is a nuclear shuttling protein, whose subcellular localization is mediated by the CRM1-dependent nuclear export pathway (PMID: 15265700). The protein is regulated transcriptionally and can be inhibited by mitochondrial proteins released in the cytoplasm upon apoptotic stimuli (PMID: 15187025). BIRC2 is also believed to be a critical regulator of vascular integrity and endothelial cell survival, thereby providing an additional target pathway for the control of angiogenesis and blood vessel homeostasis during embryogenesis, regeneration and tumorigenesis (PMID: 17934460). This BIRC2 antibody (10022-1-AP) can bind both full length (70kd) and cleaved form (60kd) of the protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Lei Wang	32882585	Biomed Pharmacother	WB
Michael Hinz	20932475	Mol Cell	WB
Dongsheng Nie	26607717	Biol Reprod	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




mouse skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 10022-1-AP (cIAP1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human testis tissue slide using 10022-1-AP (cIAP1 Antibody) at dilution of 1:200 (under 10x lens).




Immunohistochemical analysis of paraffin-embedded human testis tissue slide using 10022-1-AP (cIAP1 Antibody) at dilution of 1:200 (under 40x lens).




Immunohistochemical analysis of paraffin-embedded mouse pancreas tissue slide using 10022-1-AP (cIAP1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 10022-1-AP (cIAP1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 10022-1-AP (cIAP1 antibody), at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using cIAP1 antibody (10022-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).



1x10⁶ HepG2 cells were intracellularly stained with 0.4 ug cIAP1 Polyclonal antibody (10022-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-O-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).