

For Research Use Only

# RB1 Polyclonal antibody

Catalog Number: 10048-2-Ig

Featured Product

62 Publications



## Basic Information

**Catalog Number:**

10048-2-Ig

**Concentration:**

1200 ug/ml

**Source:**

Rabbit

**Isotype:**

IgG

**Immunogen Catalog Number:**

AG0006

**GenBank Accession Number:**

BC040540

**GeneID (NCBI):**

5925

**UNIPROT ID:**

P06400

**Full Name:**

retinoblastoma 1

**Calculated MW:**

110 kDa

**Observed MW:**

110 kDa

**Purification Method:**

Protein A purification

**Recommended Dilutions:**

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:200-1:800

## Applications

**Tested Applications:**

WB, IHC, IP, ELISA

**Cited Applications:**

WB, IHC, IF, IP, CoIP

**Species Specificity:**

human

**Cited Species:**

human, zebrafish

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

**Positive Controls:**

WB : A431 cells, Calyculin A treated Jurkat cells, MCF-7 cells

IP : A431 cells,

IHC : human intrahepatic cholangiocarcinoma tissue,

## Background Information

RB1, also named as pp110, pRb and p105 Rb, belongs to the retinoblastoma protein (RB) family. It is a key regulator of entry into cell division that acts as a tumor suppressor. RB1 acts as a transcription repressor of E2F1 target genes. The underphosphorylated, active form of RB1 interacts with E2F1 and represses its transcription activity, leading to cell cycle arrest. It is directly involved in heterochromatin formation by maintaining overall chromatin structure and, in particular, that of constitutive heterochromatin by stabilizing histone methylation. It recruits and targets histone methyltransferases SUV39H1, SUV420H1 and SUV420H2, leading to epigenetic transcriptional repression. RB1 controls histone H4 'Lys-20' trimethylation and inhibits the intrinsic kinase activity of TAF1. It mediates transcriptional repression by SMARCA4/BRG1 by recruiting a histone deacetylase (HDAC) complex to the c-FOS promoter. In resting neurons, transcription of the c-FOS promoter is inhibited by BRG1-dependent recruitment of a phospho-RB1-HDAC1 repressor complex. Upon calcium influx, RB1 is dephosphorylated by calcineurin, which leads to release of the repressor complex. In case of viral infections, interactions with SV40 large T antigen, HPV E7 protein or adenovirus E1A protein induce the disassembly of RB1-E2F1 complex thereby disrupting RB1's activity. This antibody is a rabbit polyclonal antibody raised against human RB1 fusion protein.

## Notable Publications

Author	Pubmed ID	Journal	Application
Qinghua Wang	29033588	Onco Targets Ther	WB
Shuai Huang	31660066	Theranostics	WB
Chao Zhang	36093042	iScience	WB,CoIP

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




A431 cells were subjected to SDS PAGE followed by western blot with 10048-2-Ig (RB1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




IP result of anti-RB1 (IP:10048-2-Ig, 3ug; Detection:10048-2-Ig 1:500) with A431 cells lysate 3000 ug.



Immunohistochemical analysis of paraffin-embedded human intrahepatic cholangiocarcinoma tissue slide using 10048-2-Ig (RB1 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Non-treated and Calyculin A treated Jurkat cells were subjected to SDS PAGE followed by western blot with 30376-1-AP (Phospho-RB1 (Ser807/811) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with RB1 antibody (10048-2-Ig) subsequently.



Immunohistochemical analysis of paraffin-embedded human intrahepatic cholangiocarcinoma tissue slide using 10048-2-Ig (RB1 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).