

For Research Use Only

PRC1 Polyclonal antibody

Catalog Number: 10110-2-AP

Featured Product

1 Publications



Basic Information

Catalog Number: 10110-2-AP	GenBank Accession Number: BC005140	Purification Method: Antigen affinity purification
Size: 600 µg/ml	GeneID (NCBI): 9055	Recommended Dilutions: WB 1:2000-1:12000 IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: O43663	
Isotype: IgG	Full Name: protein regulator of cytokinesis 1	
Immunogen Catalog Number: AG0163	Calculated MW: 67 kDa	
	Observed MW: 67 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : MCF-7 cells, HEK-293 cells, HeLa cells, HepG2 cells IHC : human colon cancer tissue,
Cited Applications: WB	
Species Specificity: human	
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

PRC1 is involved in cytokinesis. PRC1 is at high level during S and G2/M and drop dramatically after cell exit mitosis and enter G1. PRC1 is located in the nucleus during interphase, and becomes associated with mitotic spindles in a highly dynamic manner during mitosis, and localizes to the cell mid-body during cytokinesis. PRC1 is a good substrate for several cyclin-dependent kinases (CDKs) in vitro and is phosphorylated in vivo at sites that are phosphorylated by CDK in vitro, suggesting that PRC1 is an in vivo CDK substrate.

Notable Publications

Author	Pubmed ID	Journal	Application
Qiang Wang	37647439	Immun Inflamm Dis	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of PRC1 antibody (10110-2-AP; 1:6000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PRC1 transfected MCF-7 cells.



Various lysates were subjected to SDS PAGE followed by western blot with 10110-2-AP (PRC1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 10110-2-AP (PRC1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 10110-2-AP (PRC1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).