

For Research Use Only

# ICAM2 Polyclonal antibody

Catalog Number:10121-2-AP

Featured Product

5 Publications



## Basic Information

<b>Catalog Number:</b> 10121-2-AP	<b>GenBank Accession Number:</b> BC003097	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 250 µg/ml	<b>GeneID (NCBI):</b> 3384	<b>Recommended Dilutions:</b> WB 1:500-1:1000 IHC 1:50-1:500
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P13598	
<b>Isotype:</b> IgG	<b>Full Name:</b> intercellular adhesion molecule 2	
<b>Immunogen Catalog Number:</b> AG0165	<b>Calculated MW:</b> 31 kDa	
	<b>Observed MW:</b> 55-80 kDa	

## Applications

### Tested Applications:

IHC, WB, ELISA

### Cited Applications:

WB, IF, IHC

### Species Specificity:

human, mouse

### Cited Species:

human, mouse

### Positive Controls:

**WB** : Jurkat cells, RAW 264.7 cells, mouse brain tissue, HEK-293 cells

**IHC** : human breast cancer tissue, mouse spleen tissue, human lung cancer tissue

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

ICAM2 is a cell adhesion protein having important roles in cell migration, especially during inflammation when leukocytes cross the endothelium. Initially described as a receptor for lymphocyte function-associated antigen-1 (LFA1), ICAM2 may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell adhesion. It mediates adhesive interactions important for antigen-specific immune response, NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance. ICAM2 has six N-linked glycosylation sites at amino acids (asparagines) 47, 82, 105, 153, 178 and 187. This antibody got 55-80 kDa in western blotting maybe due to glycosylation.

## Notable Publications

Author	Pubmed ID	Journal	Application
Yoon Karina Jin KJ	18978946	PLoS One	WB,IHC
Yu Dong	24457076	Hum Pathol	IHC
Xinxin Xu	35907907	J Ovarian Res	IHC

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 10121-2-AP (ICAM2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10121-2-AP (ICAM2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10121-2-AP (ICAM2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 10121-2-AP (ICAM2 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).