

For Research Use Only

PPARD Polyclonal antibody

Catalog Number: 10156-2-AP 23 Publications



Basic Information

Catalog Number: 10156-2-AP	GenBank Accession Number: BC002715	Purification Method: Antigen affinity purification
Size: 450 µg/ml	GeneID (NCBI): 5467	Recommended Dilutions: WB 1:500-1:1000 IHC 1:10-1:100
Source: Rabbit	UNIPROT ID: Q03181	
Isotype: IgG	Full Name: peroxisome proliferator-activated receptor delta	
Immunogen Catalog Number: AG0208	Calculated MW: 50 kDa	
	Observed MW: 54 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : mouse heart tissue, mouse skeletal muscle tissue, COLO 320 cells IHC : human ovary tumor tissue,
Cited Applications: WB, IF, IHC	
Species Specificity: human, mouse, rat	
Cited Species: human, rat, mouse	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Peroxisome proliferator-activator receptor delta (PPARD), a ligand-activated transcription factor, belongs to the PPARs family, which get its name for their chemicals that induce proliferation of peroxisomes, organelles that contribute to the oxidation of fatty acids. PPARD is a receptor for peroxisome proliferator and once activated by a ligand, it binds to and activates the promoter elements of target genes, such as acylCoA oxidase gene. But it act as a repressor for the NPC1L1 gene.

Notable Publications

Author	Pubmed ID	Journal	Application
Juan Ji	36142731	Int J Mol Sci	WB
Juan Ji	27789312	Neuropharmacology	WB
Yaqiong Zhang	27815817	Neurotox Res	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse heart tissue were subjected to SDS PAGE followed by western blot with 10156-2-AP (PPAR δ antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human ovary tumor using 10156-2-AP (PPAR δ antibody) at dilution of 1:10 (under 10x lens).