For Research Use Only

S100A6 Polyclonal antibody

Catalog Number: 10245-1-AP

Featured Product

20 Publications

BC001431

6277

P06703

GeneID (NCBI):

UNIPROT ID:

Full Name:

GenBank Accession Number:



Basic Information

Catalog Number: 10245-1-AP

Size: 1000 µg/ml Source: Rabbit Isotype:

S100 calcium binding protein A6 Calculated MW:

Immunogen Catalog Number:

AG0385 10 kDa

Observed MW: 10 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:2000-1:12000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:400-1:1600 IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Cited Applications: WB, IHC, IF Species Specificity:

human **Cited Species:**

human, mouse, rat, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A549 cells, HeLa cells

IP: A549 cells,

cancer tissue

IHC: human stomach cancer tissue, human liver cancer tissue, human pancreas cancer tissue, human colon

IF/ICC: MCF-7 cells,

Background Information

S100A6 is also named as calcyclin, prolactin receptor-associated protein (PRA), growth factor-inducible protein 2A9 ir MLN4. It belongs to \$100 family of low molecular weight, acidic, calcium-binding proteins which contain two EFhand calcium binding sites. S100A6 may function as a calcium sensor to activate several processes in the calcium signal transduction pathway of cell growth, proliferation, secretion and exocytosis. The antibody is specific for S100A6.

Notable Publications

Author	Pubmed ID	Journal	Application
Anchuan Li	27596819	Mol Carcinog	WB,IHC
Elisa Teyssou	33218681	Neurobiol Aging	WB,IHC
Xu Nai-Yu NY	19492239	J Toxicol Environ Health A	WB,IHC

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




WB result of \$100A6 antibody (10245-1-AP, 1:30000) with si-Control and si-\$100A6 transfected HeLa cells..




Various lysates were subjected to SDS PAGE followed by western blot with 10245-1-AP (S100A6 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 10245-1-AP (\$100A6 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 10245-1-AP (\$100A6 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 10245-1-AP (S100A6 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 10245-1-AP (S100A6 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using \$100A6 antibody (10245-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit $\lg G(H+L)$.



1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human S100A6 (10245-1-AP) and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



IP result of anti-\$100A6 (IP:10245-1-AP, 4ug; Detection:10245-1-AP 1:5000) with A549 cells lysate 840 ug.