For Research Use Only

AURKA Polyclonal antibody

Catalog Number: 10297-1-AP

Featured Product

18 Publications

BC002499

6790

O14965 Full Name:

GeneID (NCBI):

UNIPROT ID:

GenBank Accession Number:



Purification Method:

IHC 1:100-1:400 IF/ICC 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: 10297-1-AP Concentration: 600 µg/ml

Source:
Rabbit
Isotype:
IgG
Immunogen Catalog Number:

AG0362

aurora kinase A Calculated MW: 46 kDa

Observed MW: 46 kDa

Positive Controls:

IHC: human breast cancer tissue,

IF/ICC: HeLa cells,

Applications

Tested Applications:

IHC, IF/ICC, FC (Intra), ELISA

Cited Applications:

IHC, IF

Species Specificity:

human
Cited Species:
human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

AURKA(Aurora kinase A), also named as STK6, STK15 or AIK, belongs to the Ser/Thr protein kinase family. This protein may play a role in cell cycle regulation during anaphase and/or telophase by being involved in microtubule formation and/or stabilization. Some recent evidence revealed that AURKA may also be implicated in tumor development and progression. AURKA is expressed highly in testis and various proliferating cells, migrating as a 46 kDa protein in SDS PAGE.Nuclear staining for AURKA is weak or nonexistent in normal tissue but strong in tumor tissue (PMID:19107951).

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaoman Chen	30207732	Mol Pharm	
Wei Zhou	34224810	J Ethnopharmacol	
Yanwei Liu	26468983	Oncotarget	

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using Aurora A antibody (10297-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human Aurora A (10297-1-AP) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10297-1-AP (Aurora A antibody at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10297-1-AP (Aurora A antibody at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).