For Research Use Only

PELO Polyclonal antibody

Catalog Number: 10582-1-AP

Featured Product

13 Publications



Basic Information

Catalog Number: GenBank Accession Number: 10582-1-AP BC005889 GeneID (NCBI): Concentration: 450 ug/ml 53918 Source: **UNIPROT ID:**

Q9BRX2 Rabbit Isotype: Full Name:

pelota homolog (Drosophila)

Calculated MW: Immunogen Catalog Number:

AG0873 43 kDa

> Observed MW: 43-45 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:1000-1:6000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF/ICC 1:200-1:800

Applications

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IF, IP

human, mouse, rat Cited Species:

human, mouse

Tested Applications:

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval antigen retrieval with a trace buffer and 16.0 with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, mouse testis tissue, HeLa cells,

SKOV-3 cells IP: HEK-293 cells. IHC: human liver tissue,

IF/ICC: HeLa cells,

Background Information

The Pelo gene was originally identified in a mutagenesis screen for spermatogenesis-specific genes of Drosophila melanogaster. The PELO is required for the meiotic division during the G2/M transition, and functions in recognizing stalled ribosomes and triggering endonucleolytic cleavage of the mRNA, a mechanism to release non-functional ribosomes and degrade damaged mRNAs [PMID:12556505]. It may participate in the machinery of protein synthesis or in the regulation of mRNA translation [PMID:9584085].

Notable Publications

Author	Pubmed ID	Journal	Application
Shuai Hou	31216022	Nucleic Acids Res	WB
I-Hui Wu	34107246	Cell Rep	
Mark Stoneley	35180429	Mol Cell	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various cells were subjected to SDS PAGE followed by western blot with 10582-1-AP (PELO antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




mouse testis tissue were subjected to SDS PAGE followed by western blot with 10582-1-AP (PELO antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver tissue slide using 10582-1-AP (PELO antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 10582-1-AP (PELO antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-PELO (IP:10582-1-AP, 4ug; Detection:10582-1-AP 1:500) with HEK-293 cells lysate 1200ug.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using PELO antibody (10582-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).