For Research Use Only

NUDC Polyclonal antibody

Catalog Number: 10681-1-AP

Featured Product

5 Publications



Basic Information

 Catalog Number:
 GenBank Accession Number:

 10681-1-AP
 BC006147

 Size:
 GeneID (NCBI):

 750 ug/ml
 10726

 Source:
 UNIPROT ID:

 Pabbit
 007366

Rabbit Q9Y266 Isotype: Full Name:

IgG nuclear distribution gene C homolog
(A. nidulans)

Immunogen Catalog Number: (A. nidulans)
AG0998 Calculated MW: 38 kDa

Observed MW: 42-45 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF

Species Specificity:

human
Cited Species:
human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HepG2 cells, HEK-293 cells, HeLa cells, K-562

Purification Method:

WB 1:500-1:2000

protein lysate

IHC 1:20-1:200

IF/ICC 1:200-1:800

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

cells, SH-SY5Y cells
IP: HEK-293 cells,

IHC: human pancreas cancer tissue,

IF/ICC: HeLa cells, HEK-293 cells, HepG2 cells

Background Information

Nuclear distribution protein C (NudC) is a highly conserved dynein/dynactin associated protein implicated in mitosis and cytokinesis. NudC is localized on major mitotic structures, and is involved in regulating spindle formation stability of kinetochore microtubule attachments, and chromosomes congression in early mitosis. NudC also is required for multi-protein transport along axons in neurons.

Notable Publications

Author	Pubmed ID	Journal	Application
Senyang Cao	32920902	Reprod Domest Anim	WB,IHC
Tai Young Kim	25332235	Mol Cell Biol	WB
Min Liu	34262899	Front Cell Dev Biol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 10681-1-AP (NUDC antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human pancreas cancer using 10681-1-AP (NUDC antibody) at dilution of 1:50 (under 10x lens).



IP result of anti-NUDC (IP:10681-1-AP, 4ug; Detection:10681-1-AP 1:1000) with HEK-293 cells lysate 2800ug.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using NUDC antibody (10681-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).