For Research Use Only

IFT57 Polyclonal antibody Catalog Number: 11083-1-AP Featured Product

Featured Product






Basic Information	Catalog Number: 11083-1-AP	GenBank Accession Number: BC011899		Purification Method: Antigen affinity purification	
	Concentration: GenelD (NCBI): 500 ug/ml 55081):	Recommended Dilutions: WB 1:500-1:2000	
	Source: Rabbit	UNIPROT ID: O9NWB7		WB 1.500-1.200 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500 IF/ICC 1:200-1:800	
	Isotype: IgG Immunogen Catalog Number: AG1566	Full Name:	transport 57 homolog onas)		
		Calculated M 49 kDa	Calculated MW: 49 kDa		
		Observed MW 49 kDa	<i>l</i> :		
Applications	Tested Applications: Positive Controls:			rols:	
	WB, IHC, IF/ICC, IP, ELISA		WB : HEK-293	WB : HEK-293 cells, mouse retina tissue, NIH/3T3 cells IP : HEK-293 cells,	
	Cited Applications: WB, IF, IP		IP : HEK-293 c		
				prostate cancer tissue, human pancreas	
			T-RPE1 cells,		
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	IFT57, also named as DERP8, ESRRBL1, HIPPI and MHS4R2, belongs to the IFT57 family. It is required for the formation of cilia. IFT57 plays an indirect role in sonic hedgehog signaling, cilia being required for all activity of the hedgehog pathway. It is a component of IFT complex B which composed of IFT88, IFT57, TRAF3IP1, IFT52, IFT27, HSPB11 and IFT20. In contrast to BBS proteins, IFT57 and IFT88 is found in almost every cilium.(PMID:22072986)				
Notable Publications	Author	Pubmed ID	Journal	Application	
	Malavika Raman	26389662	Nat Cell Biol	WB	
	Ying Hsu	29049287	PLoS Genet	WB	
	Laura Vuolo	30320547	Elife	WB,IF	


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of IFT57 antibody (11083-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-IFT57 transfected HEK-293 cells.




IP result of anti-IFT57 (IP:11083-1-AP, 5ug; Detection:11083-1-AP 1:800) with HEK-293 cells lysate 2000ug.



Immunohistochemical analysis of paraffinembedded human prostate cancer using 11083-1-AP (IFT57 antibody) at dilution of 1:50 (under 10x lens).



Various lysates were subjected to SDS PAGE followed by western blot with 11083-1-AP (IFT57 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed hTERT-RPE1 cells using IFT57 antibody (11083-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CoraLite®594 ARL13B antibody (CL594-17711, red).