For Research Use Only

Calretinin Polyclonal antibody

Catalog Number: 12278-1-AP 16 Publications



Basic Information

Catalog Number: 12278-1-AP

Concentration: GeneID (NCBI): 850 ug/ml 794

Observed MW: 29 kDa

BC015484

GenBank Accession Number:

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:5000-1:50000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:500-1:2000

Applications

Tested Applications: WB, IHC, IP, ELISA Cited Applications: WB, IHC, IF Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls

WB: fetal human brain tissue, human ileum tissue, mouse brain tissue, mouse colon tissue, rat brain tissue, rat colon tissue

IP: mouse brain tissue,

IHC: human appendicitis tissue, mouse brain tissue, human brain tissue, human cerebellum tissue

Background Information

Calbindin 2 (calretinin), is an intracellular calcium-binding protein belonging to the troponin C superfamily. Members of this protein family have six EF-hand domains which bind calcium. This protein plays a role in diverse cellular functions, including message targeting and intracellular calcium buffering. It also functions as a modulator of neuronal excitability, and is a diagnostic marker for some human diseases, including Hirschsprung disease and some cancers. Calretinin is a useful marker for differentiating malignant mesothelioma from carcinomas.

Notable Publications

Author	Pubmed ID	Journal	Application
Luca Pesce	34536443	Prog Biophys Mol Biol	IF
Zhihao Wang	32926169	Endocrinology	IHC
Yuki Kajita	34803625	Front Behav Neurosci	IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12278-1-AP (Calretinin antibody) at dilution of 1:12000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human appendicitis tissue slide using 12278-1-AP (Calretinin antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human appendicitis tissue slide using 12278-1-AP (Calretinin antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-Calretinin (IP:12278-1-AP, 4ug; Detection:12278-1-AP 1:10000) with mouse brain tissue lysate 1280 ug.