For Research Use Only

FBXO17 Polyclonal antibody

Catalog Number: 12844-1-AP

Featured Product

7 Publications



Basic Information

Catalog Number: GenBank Accession Number: 12844-1-AP BC012385 Concentration: GeneID (NCBI): 180 µg/ml 115290 **UNIPROT ID:** Source: Rabbit Q96EF6 Full Name: Isotype: F-box protein 17 Calculated MW: Immunogen Catalog Number:

278 aa, 31 kDa Observed MW: 29-31 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:200-1:1000 IHC 1:20-1:200 IF/ICC 1:20-1:200

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC

AG3867

Species Specificity: human, mouse Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: U-251 cells, mouse heart tissue, HepG2 cells, mouse lung tissue

IHC: human heart tissue, human prostate cancer tissue

IF/ICC: HepG2 cells, HeLa cells

Background Information

FBXO 17, also named as FBG4, FBX17, FBX26 and FBXO 26, is a substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex.

Notable Publications

Author	Pubmed ID	Journal	Application
Siyao Shen	35357059	J Gene Med	WB,IHC
Tomeka Suber	28298444	J Biol Chem	WB
Rui-Chao Chai	31157866	Carcinogenesis	IHC

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12844-1-AP (FBXO17 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




WB result of FBXO 17 antibody (12844-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-FBXO 17 transfected HepG2 cells.



Immunohistochemical analysis of paraffinembedded human heart tissue slide using 12844-1-AP (FBXO17 Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human heart tissue slide using 12844-1-AP (FBXO 17 Antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of HepG2 cells using 12844-1-AP (FBXO17 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit lpG(H+L).