## For Research Use Only

# AACS Polyclonal antibody

Catalog Number: 13815-1-AP

Featured Product

**3 Publications** 



**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 13815-1-AP
 BC040490

 Size:
 GeneID (NCBI):

 450 ug/ml
 65985

 Source:
 UNIPROT ID:

 Rabbit
 Q86V21

 Isotype:
 Full Name:

G acetoacetyl-CoA synthetase

Immunogen Catalog Number:Calculated MW:AG4899672 aa, 75 kDaObserved MW:

67 kDa, 30 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:200-1:1000 IHC 1:50-1:500 IF/ICC 1:20-1:200

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications:

WB

Species Specificity: human, mouse, rat Cited Species: mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: mouse heart tissue, mouse liver tissue

IHC: human ovary cancer tissue,

IF/ICC: MCF-7 cells,

# **Background Information**

AACS(acetoacetyl-CoA synthetase) is also named as ACSF1 and belonogs to the ATP-dependent AMP-binding enzyme family. In mammals, AACS is present in the cytosolic fractions of various tissues, especially lipogenic ones, and acetoacetate is known to be effectively incorporated into cholesterol and fatty acids. These results indicate that AACS may supply acetyl units to cytosolic compartment for cholesterol and/or fatty acid biosynthesis. It has 3 isoforms produced by alternative splicing.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
George D Dalton	34568800	iScience	WB
Faith C Kaluba	39554134	bioRxiv	WB
Eric D Queathem	39026753	bioRxiv	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



mouse heart tissue were subjected to SDS PAGE followed by western blot with 13815-1-AP (AACS Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of MCF-7 cells using 13815-1-AP (AACS antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 13815-1-AP (AACS antibody) at dilution of 1:100 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).