For Research Use Only

VAPA Polyclonal antibody

Catalog Number: 15275-1-AP

Featured Product

30 Publications



Basic Information

Catalog Number: 15275-1-AP Size: 600 ug/ml 9218 **UNIPROT ID:** Source:

Rabbit Q9PoLo Full Name: Isotype: IgG

Immunogen Catalog Number:

GenBank Accession Number:

BC002992 GeneID (NCBI):

VAMP (vesicle-associated membrane protein)-associated protein A, 33kDa

Calculated MW: 28 kDa

Observed MW: 28-30 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, CoIP Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HepG2 cells, HEK-293 cells, mouse skeletal muscle tissue, human kidney tissue, human brain tissue, mouse kidney tissue, mouse liver tissue, rat

Purification Method:

WB 1:2000-1:16000

IF/ICC 1:200-1:800

protein lysate

IHC 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

liver tissue IP: HEK-293 cells.

IHC: human colon cancer tissue, mouse heart tissue.

human liver cancer tissue

IF/ICC: HepG2 cells, HEK-293 cells

Background Information

VAPA (Vesicle-associated membrane protein-associated protein A), also known as VAP33, is a ubiquitously expressed integral membrane protein of the VAMP-associated protein (VAP) family. It is present in the plasma membrane and in intracellular vesicles, playing a role in vesicle trafficking.

Notable Publications

Author	Pubmed ID	Journal	Application
Qiaoxia Zheng	36198318	Cell	WB,CoIP
Norihiro Kotani	36152751	J Biol Chem	WB,IF
Yan G Zhao	28890335	Mol Cell	COIP


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 15275-1-AP (VAPA antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 15275-1-AP (VAPA antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 15275-1-AP (VAPA antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-VAPA (IP:15275-1-AP, 3ug; Detection:15275-1-AP 1:1000) with HEK-293 cells lysate 3200ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using VAPA antibody (15275-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using VAPA antibody (15275-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).