For Research Use Only

KPTN Polyclonal antibody

Catalog Number: 16094-1-AP

Featured Product

11 Publications



Basic Information

Catalog Number: GenBank Accession Number: 16094-1-AP BC009249
Size: GeneID (NCBI): 500 \(\mu \) g/ml 11133
Source: UNIPROT ID: Rabbit Q9Y664
Isotype: Full Name:

kaptin (actin binding protein)

Immunogen Catalog Number:Calculated MW:AG8785436 aa, 48 kDa

Observed MW: 48 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:4000 IHC 1:20-1:200

IF/ICC 1:20-1:200

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB

Species Specificity: human, mouse, rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, NIH/3T3 cells, human cerebellum tissue. mouse skeletal muscle tissue

IHC: human kidney tissue, human heart tissue, human lung tissue, human ovary tissue, human placenta tissue, human skin tissue, human spleen tissue

IF/ICC: HeLa cells,

Background Information

Kaptin, also named as KPTN and Actin-associated protein 2E4, originally isolated from human blood platelets and likely to be involved in the actin rearrangements occurring during activation. It plays a unique role in the actin rearrangements that accompany platelet activation and stereocilia formation. Kaptin is necessary for normal neuromorphogenesis. It may play a role in producing the sensory apparatus in hair cells.

Notable Publications

Author	Pubmed ID	Journal	Application
Cecilia Cattelani	34685691	Cells	WB
Kuang Shen	29590090	Nature	WB
Jose M Orozco	32719541	Nat Metab	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16094-1-AP (KPTN antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney using 16094-1-AP (KPTN antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human kidney using 16094-1-AP (KPTN antibody) at dilution of 1:100 (under 40x lens).



Immunofluorescent analysis of (-20 $^{\circ}$ Ethanol) fixed HeLa cells using 16094-1-AP (KPTN antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).