For Research Use Only

IL-12/IL-23 p40 Polyclonal antibody

Catalog Number: 17645-1-AP 4 Publications



Purification Method:

WB 1:1000-1:4000

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: 17645-1-AP Concentration: 400 ug/ml

Rabbit Isotype:

Source:

Immunogen Catalog Number:

AG11837

GenBank Accession Number:

BC067498 GeneID (NCBI): **UNIPROT ID:**

P29460

Full Name: interleukin 12B (natural killer cell

Calculated MW: 328 aa, 37 kDa Observed MW: 39 kDa

Applications

Tested Applications: WB, FC (Intra), ELISA **Cited Applications:**

WB. IF

Species Specificity:

human **Cited Species:** human, mouse

stimulatory factor 2, cytotoxic lymphocyte maturation factor 2, p40)

Positive Controls:

WB: Recombinant protein,

Background Information

IL-12 and IL-23 are heterodimeric cytokines that share a common p40 subunit (PMID: 11114383). IL-12 is composed of the IL-12 p40 subunit linked to the IL-12 p35 subunit, and the heterodimer signals through the IL-12 receptor (IL-12R), which comprises the IL-12R β 1 and IL-12R β 2 subunits. IL-23 is composed of the IL-23 p19 subunit and the IL-12 p40 (IL-12/23p40) subunit, which signals through IL-23R and IL-12R ß 1 (PMID: 11114383; 26121196). IL-12/IL-23 p40 also exists as a monomer and as a homodimer which can act as a potent IL-12 antagonist (PMID: 8958912; 18783467). IL-12/IL-23 p40 is produced by antigen-presenting cells, such as dendritic cells (DCs), monocytes, macrophages, neutrophils and, to a lesser extent, B cells (PMID: 20476918).

Notable Publications

Author	Pubmed ID	Journal	Application
Xinlong Zang	33338567	Int J Pharm	IF
Man Li	39713206	Exploration (Beijing)	WB,IF
Dwight D Harris	37940431	Surgery	


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Recombinant protein IL-12B (Eg0431), IL-12 (Eg0575), IL-12A (Ag17625) were subjected to SDS PAGE followed by western blot with 17645-1-AP (IL-12B antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours



1x10^6 Jurkat cells were intracellularly stained with 0.4 ug IL-12/IL-23 p40 Polyclonal antibody (17645-1-AP) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).