For Research Use Only

AHR Polyclonal antibody Catalog Number: 17840-1-AP Featured Produce

Featured Product






Basic Information	Catalog Number: 17840-1-AP	GenBank A BC070080	GenBank Accession Number: BC 070080		Purification Method: Antigen affinity purification	
	Concentration:GeneID (NCBI):500 µ g/ml196Source:UNIPROT ID:RabbitP35869Isotype:Full Name:IgGaryl hydrocarbon receptor		CBI):	Recommended Dilutions: WB: 1:1000-1:4000 IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC: 1:50-1:500		
			D:			
	0 0		Calculated MW: 848 aa, 96 kDa			
		Observed M 105-110 kD	served MW: 5-110 kDa			
Applications	Tested Applications: Positive Controls:					
	WB, IP, IHC, ELISA		WB: HeLa o	NB : HeLa cells, HepG2 cells, K-562 cells, MCF-7 cells		
	Cited Applications: WB, IHC, IF, CoIP, ChIP		IP : PC-3 ce	IP : PC-3 cells,		
	Species Specificity: human, mouse, rat		IHC : human lung cancer tissue, mouse small intestin tissue, human brain tissue			
	Cited Species: human, mouse, rat					
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
Background Information	The aryl hydrocarbon receptor (AhR) is a ligand-activated transcription factor that has been largely regarded as a mediator of xenobiotic metabolism [PMID:18483242]. It plays a part role in physiologic activities, including attenuation of the acute phase response, cytokine signaling, T helper (TH)17 immune cell differentiation, modulation of NF- κ B activity, and regulation of hormonal signaling [PMID:20423157,18540824]. It also mediates transcription factor sequestering away from a gene promoter or tethering of the AhR to a transcription factor on a promoter. AHR calculated molecular masses differ by <10%, compared with the apparent molecular masses predicted from SDS-PAGE for the two receptors (105 and 95 kDa, respectively). (PMID: 8246913)					
Notable Publications	Author	Pubmed ID	Journal		Application	
	Susu Lin	36169685	Psychopharmacolog	y (Berl)	WB	
	Rongyao Liang	34555451	J Ethnopharmacol		WB	
	Jie Xiong	32941808	Exp Cell Res		WB	
Storage	Storage: Store at -20°C. Stable for one ye Storage Buffer: PBS with 0.02% sodium azide a Aliquoting is unnecessary for -2	nd 50% glycerol, ۱				


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 17840-1-AP (AHR antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 17840-1-AP (AHR antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



IP result of anti-AHR (IP:17840-1-AP, 4ug; Detection:17840-1-AP 1:500) with PC-3 cells lysate 1880ug.



WB result of AHR antibody (17840-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-AHR transfected HepG2 cells.