## For Research Use Only

## NF-M-Specific Polyclonal antibody

Catalog Number: 20664-1-AP

2 Publications



**Basic Information** 

Catalog Number: 20664-1-AP Size: 700 µg/ml

Source: UNIPROT ID:
Rabbit P07197
Isotype: Full Name: neurofilame

neurofilament, medium polypeptide Calculated MW:

102 kDa

Observed MW:
140-160 kDa

NM\_005382

4741

GeneID (NCBI):

GenBank Accession Number:

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF/ICC 1:50-1:500

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Cited Applications:

WB

Species Specificity: human, mouse, rat Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls

WB: mouse brain tissue, rat brain tissue, human brain

tissue

IP: rat brain tissue,

IHC: human brain tissue, rat brain tissue

IF/ICC: SH-SY5Y cells,

**Background Information** 

NEFM, also named as NEF3 and NFM, belongs to the intermediate filament family. Neurofilaments are the 10nm intermediate filaments found specifically in neurons. They are a major component of the cell's cytoskeleton, and provide support for normal axonal radial growth. Neurofilaments usually contain three intermediate filament proteins: L, M, and H which are involved in the maintenance of neuronal caliber. The names given to the three major neurofilament subunits are based upon the apparent molecular weight of the mammalian subunits on SDS-PAGE: NF-L, 65-68 kDa; NF-M,145-160 kDa and NF-H, 200-220 kDa. The antibody is specific to NEFM.

**Notable Publications** 

Author	Pubmed ID	Journal	Application
Peng-Peng Zhu	35348668	Hum Mol Genet	WB
Markus T Sainio	35237613	Front Cell Dev Biol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

torage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data




mouse brain and rat brain tissues were subjected to SDS PAGE followed by western blot with 20664-1-AP (NF-M-Specific antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




WB result of 20664-1-AP.




IP result of anti-NF-M-Specific (IP:20664-1-AP, 4ug; Detection:20664-1-AP 1:1000) with rat brain tissue lysate 4000ug.




Immunohistochemical analysis of paraffinembedded human brain tissue slide using 2064-1-AP (NF-M-Specific antibody at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 20664-1-AP (NF-M-Specific antibody at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (-20° Ethanol) fixed SH-SY5Y cells using 20664-1-AP (NF-M-Specific antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 PC-12 cells were intracellularly stained with 0.4 ug Anti-Human NF-M-Specific (20664-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).