For Research Use Only

OLIG2 Polyclonal antibody

Catalog Number:25754-1-AP



Basic Information	Catalog Number: 25754-1-AP	GenBank Accession Numb BC047511	er: Purification Method: Antigen affinity purification	
	Size: 750 ug/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG22642	GenelD (NCBI): 10215	Recommended Dilutions: WB 1:500-1:1000	
		UNIPROT ID: Q13516	IHC 1:50-1:500 IF-P 1:50-1:500	
		Full Name: oligodendrocyte lineage transcription factor 2 Calculated MW: 32 kDa		
				Observed MW: 40 kDa
		Applications		
WB, IHC, IF-P, ELISA	WE		WB : rat brain tissue, C6 cells	
Species Specificity: human, mouse, rat			C : human gliomas tissue, mouse cerebellum tissue, puse brain tissue	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			P : mouse brain tissue,	
Background Information	The oligodendrocyte lineage-specific basic helix-loop-helix (OLIG) family of transcription factors include OLIG1- OLIG3, which differ in tissue expression. OLIG1 and OLIG2 are specifically expressed in nervous tissue as gene regulators of oligodendrogenesis. OLIG1 and OLIG2 interact with the Nkx-2.2 homeodomain protein, which is responsible for directing ventral neuronal patterning in response to graded Sonic hedgehog signaling in the embryonic neural tube. These interactions between OLIG proteins and Nkx-2.2 appear to promote the formation of alternate cell types by inhibiting V3 interneuron development. OLIG1 and OLIG2 are abundantly expressed in oligodendroglioma and nearly absent in astrocytomas. Therefore, OLIG proteins are candidates for molecular markers of human glial brain tumors, which are the most common primary malignancies of the human brain.			
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and	•		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 25754-1-AP (OLIG2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



rat brain tissue were subjected to SDS PAGE followed by western blot with 25754-1-AP (OLIG2 Antibody) at dilution of 1:600 incubated at 4 degree celsius over night.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using OLIG2 antibody (25754-1-AP) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using OLIG2 antibody (25754-1-AP) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).