

For Research Use Only

GTF3C2 Polyclonal antibody

Catalog Number: 27494-1-AP

1 Publications



Basic Information

Catalog Number:

27494-1-AP

Size:

300 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG26666

GenBank Accession Number:

BC020981

GeneID (NCBI):

2976

UNIPROT ID:

Q8WUA4

Full Name:

general transcription factor IIIC,
polypeptide 2, beta 110kDa

Calculated MW:

101 kDa

Observed MW:

100-110 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IHC 1:50-1:500

IF/ICC 1:400-1:1600

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB

Species Specificity:

human

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293T cells, A549 cells, HeLa cells, HepG2 cells, MCF-7 cells, PC-3 cells

IHC: human thyroid cancer tissue, human cervical cancer tissue

IF/ICC: HeLa cells,

Background Information

GTF3C2, also named as KIAA0011, is a 911 amino acid protein, which localizes in the nucleus. GTF3C2 is required for RNA polymerase III-mediated transcription. GTF3C2 is a component of TFIIC that initiates transcription complex assembly on tRNA and is required for transcription of 5S rRNA and other stable nuclear and cytoplasmic RNAs. GTF3C2 may play a direct role in stabilizing interactions of TFIIC2 with TFIIC1.

Notable Publications

Author	Pubmed ID	Journal	Application
Aiko Iwata-Otsubo	38520561	Hum Genet	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 27494-1-AP (GTF3C2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human thyroid cancer tissue slide using 27494-1-AP (GTF3C2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using GTF3C2 antibody (27494-1-AP) at dilution of 1:800 and Multi-rAb CoraLite @ Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).