

For Research Use Only

# BCL2 Monoclonal antibody

Catalog Number: 60178-1-Ig

Featured Product

315 Publications



## Basic Information

**Catalog Number:**

60178-1-Ig

**Concentration:**

1500 ug/ml

**Source:**

Mouse

**Isotype:**

IgG2b

**Immunogen Catalog Number:**

AG3508

**GenBank Accession Number:**

BC027258

**GeneID (NCBI):**

596

**UNIPROT ID:**

P10415

**Full Name:**

B-cell CLL/lymphoma 2

**Calculated MW:**

26 kDa

**Observed MW:**

26 kDa

**Purification Method:**

Protein A purification

**CloneNo.:**

4H8C6

**Recommended Dilutions:**

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:6400-1:25600

IF-P 1:50-1:500

IF/ICC 1:200-1:800

## Applications

**Tested Applications:**

IHC, IF/ICC, IF-P, IP, ELISA

**Cited Applications:**

IHC, IF, IP, CoIP

**Species Specificity:**

human, pig

**Cited Species:**

human, pig, rabbit, monkey, bovine

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

**Positive Controls:**

IP: U-937 cells,

IHC: human appendicitis tissue, human lymphoma tissue, human tonsillitis tissue, Jurkat cells

IF-P: human appendicitis tissue, human tonsillitis tissue

IF/ICC: MCF-7 cells,

## Background Information

BCL2 belongs to the Bcl-2 family. It suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural cells. BCL2 regulates cell death by controlling the mitochondrial membrane permeability. It appears to function in a feedback loop system with caspases. BCL2 inhibits caspase activity either by preventing the release of cytochrome c from the mitochondria and/or by binding to the apoptosis-activating factor (APAF-1).

## Notable Publications

Author	Pubmed ID	Journal	Application
Ji Xing	36230734	Cancers (Basel)	IF
Fei Yu	36175965	Chin Med	WB
Jinliang Liu	34630847	Oxid Med Cell Longev	WB

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human appendicitis tissue slide using 60178-1-Ig (BCL2 antibody) at dilution of 1:12800 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human appendicitis tissue slide using 60178-1-Ig (BCL2 antibody) at dilution of 1:12800 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using BCL2 antibody (60178-1-Ig, Clone: 4H8C6) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



IP result of anti-BCL2 (IP:60178-1-Ig, 5ug; Detection:60178-1-Ig 1:1000) with U-937 cells lysate 4000ug.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 60178-1-Ig (BCL2 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human appendicitis tissue using BCL2 antibody (60178-1-Ig, Clone: 4H8C6) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).