### For Research Use Only

# P53 Monoclonal antibody

Catalog Number:60283-2-lg

Featured Product

278 Publications

BC003596

GeneID (NCBI):

**UNIPROT ID:** 

Full Name:

tumor protein p53

P04637

GenBank Accession Number:



**Basic Information** 

Catalog Number: 60283-2-lg Concentration:

1000 ug/ml Source:

Mouse Isotype: lgG2b

Immunogen Catalog Number:

AG0698

Calculated MW: 44 kDa

> Observed MW: 53 kDa

**Purification Method:** 

Protein A purification

CloneNo.: 6C4B6

Recommended Dilutions: WB 1:5000-1:50000

IHC 1:1600-1:6400 IF/ICC 1:400-1:1600

# **Applications**

**Tested Applications:** WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC, IF, IP, CoIP

Species Specificity: human, rat

**Cited Species:** 

human, rat, pig, rabbit, canine, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB: A431 cells, COLO 320 cells, HSC-T6 cells, ROS1728 cells, rat thymus tissue, HaCaT cells, HT-29 cells, HEK-293 cells

IHC: human colon cancer tissue, human breast cancer tissue, human gliomas tissue, human lung cancer tissue, human ovary cancer tissue, human ovary tumor tissue, human stomach cancer tissue, human tonsillitis

IF/ICC: A431 cells.

# **Background Information**

TP53, also known as P53 and NY-CO-13, belongs to the p53 family and has 9 isoforms. In SDS-Page, the observed molecular weight is about 53 kDa. TP53 acts as a tumor suppressor in many tumor types, including growth arrest or apoptosis depending on the physiological circumstances and cell types. It is involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. TP53 Localizes in the nucleus in most cells but found in the cytoplasm in some cells. (PMID: 26166714; PMID: 25225161)

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Jie Liu	36238648	Oxid Med Cell Longev	WB
Xiaofang Zhu	36174712	Life Sci	WB
Yingjie Qing	34603598	Oxid Med Cell Longev	WB

# Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


#### Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 60283-2-1g (P53 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 60283-2-lg (P53 antibody) at dilution of 1:3200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




HSC-T6 cells were subjected to SDS PAGE followed by western blot with 60283-2-Ig (P53 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




ROS1728 cells were subjected to SDS PAGE followed by western blot with 60283-2-lg (P53 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



rat thymus tissue were subjected to SDS PAGE followed by western blot with 60283-2-lg (P53 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 60283-2-lg (P53 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using P53 antibody (60283-2-1g, Clone: 6C4B6) at dilution of 1:800 and Coralite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-phalloidin (red).