For Research Use Only

IL-2 Monoclonal antibody

Catalog Number: 60306-1-lg 10 Publications



Basic Information

Applications

Catalog Number: 60306-1-lg

Concentration: 2600 ug/ml Source: Mouse Isotype: IgG2a

Immunogen Catalog Number:

AG10199

GenBank Accession Number:

BC070338 GeneID (NCBI): 3558 UNIPROT ID: P60568 Full Name: interleukin 2 Calculated MW:

153 aa, 18 kDa Observed MW:

18 kDa

Tested Applications:

WB, IHC, IF/ICC, IF-P, ELISA Cited Applications:

WB, IHC

Species Specificity:

human **Cited Species:**

human, mouse, pig, hamster

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, MOLT-4 cells, human peripheral blood leukocyte

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:1000-1:4000 IHC 1:20-1:200

IF-P 1:200-1:800

IF/ICC 1:400-1:1600

CloneNo.:

2F11C5

IHC: human tonsillitis tissue, human breast cancer

tissue

IF-P: human tonsillitis tissue.

IF/ICC: Jurkat cells,

Background Information

IL2 is a secreted cytokine involved in the operation and regulation of immune system. IL2 plays a pivotal role in Tcell activation and effector functions, including T cell proliferation, interferon- γ production and cytotoxicity. IL2 can stimulate the propagation of lymphocytes and induce the cytotoxic T lymphocytes and lymphokine-activated killer cells for tumor cells. IL2 also influences memory T cell homeostasis through the regulation of memory T cell numbers and enhances major histocompatibility complex II (MHC II) expression on tumor cells.

Notable Publications

Author	Pubmed ID	Journal	Application
Chengfen Wang	25264893	PLoS One	WB
Hai long Yang	34657991	Inflammation	WB
Yinrui Guo	34045460	Transl Psychiatry	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 60306-1-1g (IL-2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60306-1-lg (IL2 Antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieved with Tris-EDTA buffer, pH9.0.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60306-1-lg (IL2 Antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieved with Tris-EDTA buffer, pH9.0.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human tonsillitis tissue using IL-2 antibody (60306-1-Ig, Clone: 2F11C5) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using IL-2 antibody (60306-1-Ig, Clone: 2F11C5) at dilution of 1:800 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1).