#### For Research Use Only

# c-SRC Monoclonal antibody

Catalog Number: 60315-1-lg 16 Publications



**Basic Information** 

Catalog Number: 60315-1-lg Size:

1500 ug/ml Source: Mouse Isotype: lgG2b

Immunogen Catalog Number:

AG22306

GenBank Accession Number:

BC011566 GeneID (NCBI): 6714

ENSEMBL Gene ID: ENSG00000197122 **UNIPROT ID:** P12931 Full Name:

v-src sarcoma (Schmidt-Ruppin A-2) viral oncogene homolog (avian)

Calculated MW: 60 kDa Observed MW: 60 kDa

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, ELISA Cited Applications:

Species Specificity: human, mouse, rat, hamster

**Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, RAW 264.7 cells, CHO cells, HepG2 cells, HEK-293 cells, Jurkat cells, NIH/3T3 cells, 4T1 cells, PC-12 cells, HSC-T6 cells

**Purification Method:** 

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:100-1:400

IF/ICC 1:50-1:500

CloneNo.:

5E10C4

IHC: human breast cancer tissue,

IF/ICC: MCF-7 cells,

## **Background Information**

SRC, also named as SRC1 and p60-Src, belongs to the protein kinase superfamily, Tyr protein kinase family and SRC  $subfamily. \ It is a non-receptor protein tyrosine kinase that plays pivotal \ roles in numerous cellular processes such a contract of the plays pivotal roles in numerous cellular processes such a contract of the plays pivotal roles in numerous cellular processes and the pl$ as proliferation, migration, and transformation. In concert with PTK2B, SRC plays an important role in osteoclastic  $bone\ resorption.\ It\ promotes\ energy\ production\ in\ osteoclasts\ by\ activating\ mitochondrial\ cytochrome\ C\ oxidase.$ 

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Fan Yang	26347956	Nat Commun	WB
Márcia Faria	36358781	Cancers (Basel)	WB
Shiyu Wu	34821070	Adv Sci (Weinh)	IF

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 60315-1-1g (c-SRC antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60315-1-Ig (c-SRC Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60315-1-Ig (c-SRC Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 60315-1-1g (c-SRC antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using c-SRC antibody (60315-1-lg, Clone: 5E10C4) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).