## For Research Use Only

## ACTN4 Monoclonal antibody


Catalog Number:66628-1-lg Featured Product 4 Publications



Basic Information	Catalog Number: 66628-1-lg	GenBank Ad BC005033	cession Number:	Purification Method: Protein G purification	
	Concentration: 1000 ug/ml			CloneNo.: 1E12B8	
	Source: Mouse	UNIPROT IE O43707	):	Recommended Dilutions: WB 1:5000-1:20000 IHC 1:2500-1:10000	
	lsotype: lgG1	Full Name: actinin, alp	ha 4		
	Immunogen Catalog Number: AG13549	Calculated 105 kDa	MW:		
		Observed N 105 and 80			
Applications	Tested Applications: WB, IHC, ELISA				
	Cited Applications: WB, IF, IP	cells, Jurkat	WB: HeLa cells, LNCaP cells, HEK-293 cells, MCF-7 cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells, 4T1 cells, K-562 cells, pig brain tissue, mouse brain tissu		
	Species Specificity: rat brain tissue, ch			brain tissue, chicken brain tissue	
				i bleast cancer tissue, numan cun	ig cancer
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	ACTN4 (alpha actinin 4) is a ubiquitous actin-binding protein that cross-links actin filaments into bundles to form filopodia, and has a crucial role in cytoskeletal organization. ACTN4 is colocalized with actin stress fibers and is dispersed in the cytoplasm and in the nucleus. Increased levels of ACTN4 have been associated with cancer metastasis, indicating its implication in cancer cell migration and metastasis. Defects in ACTN4 cause glomerular disease. This antibody recognizes 105 and 80 kDa of endogenous ACTN4 protein in various lysates. (22370640, 10700177)				
Notable Publications	Author	Pubmed ID	Journal	Applica	tion
	Bing Yu	34758440	Int Immunopharmac	ol IF	
	Xiaoying Liu	34061428	Environ Toxicol W		
	Jianlin Zhao	31399951	J Physiol Biochem	WB,IF	
Storage	Storage: Store at -20°C. Stable for one ye Storage Buffer:	ear after shipment.			

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 66628-1-lg (ACTN4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66628-1-1g (ACTN4 antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).