For Research Use Only

ASF/SF2 Monoclonal antibody

Catalog Number:66671-1-lg 2 Publications



Basic Information

Catalog Number: 66671-1-lg Size:

1000 µg/ml Source: Mouse Isotype:

Immunogen Catalog Number:

AG3626

lgG1

GenBank Accession Number: BC010264

GeneID (NCBI): 6426 **UNIPROT ID:** Q07955

Full Name: splicing factor, arginine/serine-rich 1 protein lysate Calculated MW:

Observed MW: 32 kDa, 28 kDa, 22 kDa

248 aa, 28 kDa

Positive Controls:

WB: HeLa cells, LNCaP cells, HEK-293 cells, HepG2 cells, Jurkat cells, HSC-T6 cells, PC-12 cells, NIH/3T3

Purification Method:

Protein G purification

Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

WB 1:5000-1:50000

IHC 1:200-1:800

IF/ICC 1:200-1:800

CloneNo.:

1A6C5

cells, RAW 264.7 cells IP: HeLa cells,

IHC: human ovary tumor tissue, human gliomas tissue

IF/ICC: HepG2 cells,

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

IF, IP

Species Specificity: Human, mouse, rat Cited Species: human, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

SFRS1, also named as ASF, SF2, SF2P33, SFRS1 and ASF-1, belongs to the splicing factor SR family. It plays a role in preventing exon skipping, ensuring the accuracy of splicing and regulating alternative splicing. SFRS1 interacts with other spliceosomal components, via the RS domains, to form a bridge between the 5'- and 3'-splice site binding components, U1 snRNP and U2AF. Isoform ASF-2 and isoform ASF-3 act as splicing repressors. This antibody can recognize all the 3 isoforms(28kd,33kd and 22kd) of SFRS1.

Notable Publications

Author	Pubmed ID	Journal	Application
Lu Li	39207136	J Virol	IF
Hannah C Happ	37205386	bioRxiv	IP

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66671-1-1g (ASF/SF2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



IP result of anti-ASF/SF2 (IP:66671-1-Ig, 5ug; Detection:66671-1-Ig 1:5000) with HeLa cells lysate 2200 ug.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 66671-1-1g (ASF/SF2 antibody) at dilution of 1:400 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using ASF/SF2 antibody (66671-1-lg, Clone: 1A6C5) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).