

For Research Use Only

# CHOP/GADD153 Monoclonal antibody

Catalog Number: 66741-1-Ig **43 Publications**



## Basic Information

<b>Catalog Number:</b> 66741-1-Ig	<b>GenBank Accession Number:</b> BC003637	<b>Purification Method:</b> Protein A purification
<b>Concentration:</b> 3400 ug/ml	<b>GeneID (NCBI):</b> 1649	<b>CloneNo.:</b> 4F3G1
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> P35638	<b>Recommended Dilutions:</b> WB 1:1000-1:6000 IHC 1:250-1:1000
<b>Isotype:</b> IgG2a	<b>Full Name:</b> DNA-damage-inducible transcript 3	
<b>Immunogen Catalog Number:</b> AG7354	<b>Calculated MW:</b> 19 kDa	
	<b>Observed MW:</b> 30 kDa	

## Applications

<b>Tested Applications:</b> WB, IHC, ELISA	<b>Positive Controls:</b> WB : HSC-T6 cells, HepG2 cells, C6 cells, NIH/3T3 cells, Tunicamycin treated HepG2 cells
<b>Cited Applications:</b> WB, IHC, IF	<b>IHC :</b> human cervical cancer tissue, human skin cancer tissue
<b>Species Specificity:</b> human, mouse, rat	
<b>Cited Species:</b> human, mouse, rat, pig, rabbit, shrew	

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

CHOP, also known as GADD153 or DDIT3, is a highly conserved gene in both the structural and regulatory regions. Imposed by unfolded and misfolded proteins, CHOP is significantly induced by ER stress. CHOP is considered a proapoptotic marker of ER stress dependent cell death. CHOP acts as a dominant-negative inhibitor of the transcription factor C/EBP and LAP. It may play an important role in the malignant transformation of nevus to melanoma. The calculated molecular weight of CHOP is 19 kDa, but the protein migrates on an SDS-PAGE gel with an observed molecular mass of 29 kDa (PMID: 1547942).

## Notable Publications

Author	Pubmed ID	Journal	Application
Qi Xu	36341965	Environ Toxicol Pharmacol	WB
Yujie Zhong	36501024	Nutrients	WB
Xiao Zheng	34748795	Dev Comp Immunol	WB


## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol, pH7.3  
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:  
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66741-1-Ig (CHOP; GADD153 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Un-treated and Tunicamycin treated HepG2 lysates were subjected to SDS PAGE followed by western blot with 66741-1-Ig (CHOP; GADD153 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 66741-1-Ig (CHOP; GADD153 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 66741-1-Ig (CHOP; GADD153 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).