

XRN2 Monoclonal antibody

Catalog Number: 66852-1-Ig 2 Publications

Basic Information

Catalog Number: 66852-1-Ig	GenBank Accession Number: BC006417	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 22803	CloneNo.: 2C3E3
Source: Mouse	UNIPROT ID: Q9H0D6	Recommended Dilutions: WB 1:1000-1:6000 IHC 1:150-1:600 IF/ICC 1:400-1:1600
Isotype: IgG1	Full Name: 5'-3' exoribonuclease 2	
Immunogen Catalog Number: AG27927	Calculated MW: 104 kDa	
	Observed MW: 109 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, FC (Intra), ELISA	Positive Controls: WB : HT-29 cells, HEK-293 cells, COLO 320 cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells IHC : human breast cancer tissue, IF/ICC : MCF-7 cells,
Cited Applications: WB	
Species Specificity: human, mouse, rat	
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

XRN2 is one exonuclease that degrades the Pol II associated product of poly(A) site cleavage, which is crucial for Pol II termination. During transcription termination, XRN2 cleaves at the polyadenylation site liberates a 5' fragment which is subsequently processed to form the mature mRNA and a 3' fragment which remains attached to the elongating polymerase. The processive degradation of this 3' fragment by this protein may promote termination of transcription.

Notable Publications

Author	Pubmed ID	Journal	Application
Wen-Long Xue	32186933	Am J Physiol Cell Physiol	WB
Ruihui Xie	36939377	Cancer Res	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66852-1-Ig (XRN2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66852-1-Ig (XRN2 antibody) at dilution of 1:300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using XRN2 antibody (66852-1-Ig, Clone: 2C3E3) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-phalloidin (red).



1×10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human XRN2 (66852-1-Ig, Clone:2C3E3) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).