## For Research Use Only

## ABCA7 Monoclonal antibody

Catalog Number: 67128-1-lg



**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 67128-1-lg
 NM\_019112

 Concentration:
 GeneID (NCBI):

 1000 ug/ml
 10347

 Source:
 UNIPROT ID:

 Mouse
 Q8IZY2

Isotype: Full Name:
IgG1 ATP-binding cassette, sub-family A
Immunogen Catalog Number: (ABC1), member 7

AG27499 Calculated MW: 234 kDa
Observed MW:

250 kDa

**Applications** 

Tested Applications: WB, IHC, IF-P, ELISA Species Specificity: human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: 37°C incubated rat brain tissue, Jurkat cells, mouse brain tissue, pig brain tissue, pig thymus tissue, unboiled mouse brain tissue, unboiled rat brain tissue

**Purification Method:** 

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:50-1:500

IF-P 1:200-1:800

CloneNo.:

2E1A10

IHC: human thymus tissue,
IF-P: human tonsillitis tissue,

## **Background Information**

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Immunohistochemical analysis of paraffinembedded human thymus tissue slide using 67128-1-Ig (ABCA7 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using ABCA7 antibody (67128-1-Ig, Clone: 2E1A10) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using ABCA7 antibody (67128-1-lg, Clone: 2E1A10) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 67128-1-lg (ABCA7 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.