For Research Use Only

ACLY Monoclonal antibody

Catalog Number: 67166-1-lg 8 Publications



Basic Information

Catalog Number: 67166-1-Ig Concentration:

1000 µ g/ml
Source:
Mouse
Isotype:
IgM

Immunogen Catalog Number: AG7709

Observed MW:

120 kDa

121 kDa

BC006195

GeneID (NCBI):

UNIPROT ID:

Full Name:

ATP citrate lyase Calculated MW:

P53396

GenBank Accession Number:

Purification Method:

Protein A purification

CloneNo.: 1E10D5

Recommended Dilutions: WB 1:1000-1:10000 IHC 1:1000-1:4000 IF/ICC 1:400-1:1600

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications:

WB, IF, IP Species Specificity: Human, Mouse, Rat Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: Jurkat cells, MCF-7 cells, HSC-T6 cells

IHC: human prostate cancer tissue,

IF/ICC: A549 cells,

Background Information

ACLY(ATP-citrate synthase) is also named as ACL It is the primary enzyme responsible for the synthesis of cytosolic acetyl-CoA. ACLY serves as not only a target in oxygenated cells for suppression of lipid synthesis and histone acetylation, but also as a susceptible target in hypoxic cells to restore inhibition of glycolysis. In nonsmall cell lung carcinoma and hepatocellular carcinoma, ACLY is overexpressed compared with normal parenchyma suggesting that ACLY may represent a common target among highly malignant tumors(PMID:19795461). This protein has 2 isoforms produced by alternative splicing.

Notable Publications

Author	Pubmed ID	Journal	Application
Yu Hu	34539243	Mediators Inflamm	WB
Feifei Zhou	36241648	Cell Death Dis	WB,IF,IP
Ruixin Wu	39798341	Phytomedicine	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 67166-1-Ig (ACLY antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 67166-1-Ig (ACLY antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using ACLY antibody (67166-1-lg, Clone: 1£10D5) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).