For Research Use Only

Phospho-Marcksl1 Polyclonal antibody



Catalog Number: 10018-3-AP

5 Publications

Basic Information

Catalog Number: 10018-3-AP Size: 392 µ g/ml Source: Rabbit Isotype: IgG GenBank Accession Number:

BCO46601 GeneID (NCBI): 17118 UNIPROT ID: P26645 Full Name:

myristoylated alanine rich protein

kinase C substrate Calculated MW: 32 kDa Observed MW: 32 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:3000

Applications

Tested Applications: WB, ELISA Cited Applications: WB

Species Specificity: human, mouse Cited Species: bovine, human, rat

Positive Controls:

WB: Mouse J774 Macrophage cellss-treated with PMA, HeLa cells, Mouse J774 Macrophage cells

Background Information

MARCKS is a member of MARCKS family of PKC substrate and it is also a widely accepted indicator of PKC activation. It shows a ubiquitous tissue distribution. The calculated molecular weight of MARCKS is 39 kDa and its apparent molecular weights were found at 68 kDa to 90 kDa in different species. This protein is a known to regulated cytoskeleton remodeling and membrane recycling. This antibody is raised against phospho-Marcksl1 (genel D:17357) with S93/S104 phosphorylated. Because of the sequence similarity, this antibody can recognize phospho-Marcksl1 and phospho-Marcks.

Notable Publications

Author	Pubmed ID	Journal	Application
Fiala Milan M	19440461	Open Virol J	WB
Peng Y-J YJ	19369559	J Neurosci	WB
Jin Benjamin Y BY	22927426	Proc Natl Acad Sci U S A	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data





WB result of phospho-Marcks antibody (10018-3-AP, 1:1500) with Mouse J774 Macrophage cells-treated with PMA.

HeLa cells were subjected to SDS PAGE followed by western blot with 10018-3-AP (Marcks Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.