For Research Use Only

ERBB3 Polyclonal antibody

Catalog Number:10369-1-AP 9 Publications





Basic Information	Catalog Number: 10369-1-AP	GenBank Accession Number: BC002706	Purification Method: Antigen affinity purification	
	Size:	GeneID (NCBI):	Recommended Dilutions:	
	500 µg/ml	2065	IHC 1:50-1:500	
	Source: Rabbit	UNIPROT ID: P21860		
	Isotype: IgG Immunogen Catalog Number: AG0215	Full Name: v-erb-b2 erythroblastic leukemia		
		viral oncogene homolog 3 (avian)		
		Calculated MW: 148 kDa		
Applications	Tested Applications:	Positive Controls:		
	IHC, ELISA Cited Applications: WB, IHC		IHC : human breast cancer tissue, human prostate cancer tissue	
	Species Specificity: human			
	Cited Species: human, mouse			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
	V-erb-b2 erythroblastic leukemia viral oncogene homolog 3 (ErbB-3, HER3) is a member of the EGF receptor tyrosi kinase family including EGFR (HER1), Neu (ErbB-2, HER2), ErbB-3 (HER3), and ErbB-4 (HER4) that are frequently overexpressed in a variety of carcinomas. These family members form either homodimers or heterodimers upon ligand binding to mediate cell growth. ErbB-3 binds and is activated by neuregulins and NTAK. It forms a heterodimer with each of the other ErbB receptors. ErbB-3 is predominantly expressed in epithelial tissues and bra and is overexpressed in a subset of human mammary tumors.			
Background Information	kinase family including EGFR (HE overexpressed in a variety of carc ligand binding to mediate cell gro heterodimer with each of the othe	R1), Neu (ErbB-2, HER2), ErbB-3 (HER3 inomas. These family members form owth. ErbB-3 binds and is activated by r ErbB receptors. ErbB-3 is predomina), and ErbB-4 (HER4) that are frequently either homodimers or heterodimers upon neuregulins and NTAK. It forms a	
	kinase family including EGFR (HE overexpressed in a variety of carc ligand binding to mediate cell gro heterodimer with each of the othe and is overexpressed in a subset o	R1), Neu (ErbB-2, HER2), ErbB-3 (HER3 inomas. These family members form owth. ErbB-3 binds and is activated by r ErbB receptors. ErbB-3 is predomina), and ErbB-4 (HER4) that are frequently either homodimers or heterodimers upon neuregulins and NTAK. It forms a	
	kinase fam ⁱ ly including EGFR (HE overexpressed in a variety of carc ligand binding to mediate cell gro heterodimer with each of the othe and is overexpressed in a subset o	R1), Neu (ErbB-2, HER2), ErbB-3 (HER3 inomas. These family members form owth. ErbB-3 binds and is activated by r ErbB receptors. ErbB-3 is predomina of human mammary tumors.), and ErbB-4 (HER4) that are frequently either homodimers or heterodimers upon neuregulins and NTAK. It forms a ntly expressed in epithelial tissues and bra Application	
	kinase family including EGFR (HE overexpressed in a variety of carc ligand binding to mediate cell gro heterodimer with each of the othe and is overexpressed in a subset of Author Hongbing Li	R1), Neu (ErbB-2, HER2), ErbB-3 (HER3 inomas. These family members form owth. ErbB-3 binds and is activated by rr ErbB receptors. ErbB-3 is predomina of human mammary tumors. Pubmed ID Journal), and ErbB-4 (HER4) that are frequently either homodimers or heterodimers upon neuregulins and NTAK. It forms a ntly expressed in epithelial tissues and bra Application	
Background Information	kinase family including EGFR (HE overexpressed in a variety of carc ligand binding to mediate cell gro heterodimer with each of the othe and is overexpressed in a subset of Author Hongbing Li Samusi Adediran	R1), Neu (ErbB-2, HER2), ErbB-3 (HER3) inomas. These family members form owth. ErbB-3 binds and is activated by r ErbB receptors. ErbB-3 is predomina of human mammary tumors. Pubmed ID Journal 31556771 Acta Otolaryngo), and ErbB-4 (HER4) that are frequently either homodimers or heterodimers upon neuregulins and NTAK. It forms a ntly expressed in epithelial tissues and bra Application IHC	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10369-1-AP (ERBB3 Antibody) at dilution of 1:200 (under 10x lens). Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10369-1-AP (ERBB3 Antibody) at dilution of 1:200 (under 40x lens).