For Research Use Only

CDK5 Polyclonal antibody

Catalog Number: 10430-1-AP

Featured Product

12 Publications



Basic Information

Catalog Number: GenBank Accession Number: 10430-1-AP BC005115

Size: GeneID (NCBI): 1800 \(\mu \) g/ml 1020

Source: UNIPROT ID: Rabbit Q00535

Isotype: Full Name:

gG cyclin-dependent kinase 5

Immunogen Catalog Number:Calculated MW:AG068631 kDa, 33 kDaObserved MW:

33 kDa

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA Cited Applications: WB

VVD

Species Specificity: human, mouse, rat Cited Species: human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HepG2 cells, mouse pancreas tissue, NIH/3T3

Purification Method:

WB 1:500-1:2000 IHC 1:50-1:500

IF 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

cells, mouse brain tissue

IHC: mouse kidney tissue, human renal cell carcinoma

tissue

IF: SH-SY5Y cells,

Background Information

Cyclin-dependent kinase 5 (CDK5), belongs to the cyclin-dependent kinase family, is a proline-directed serine/threonine-protein kinase that essential for neuronal cell cycle arrest and differentiation and may be involved in apoptotic cell death in neuronal diseases by triggering abortive cell cycle re-entry. CDK5 predominantly expressed in neurons where it phosphorylates both high molecular weight neurofilaments and microtubule-associated protein tau.

Notable Publications

Author	Pubmed ID	Journal	Application
Yue Li	29031852	Neuropharmacology	WB
Liu Wei W	22614921	J Cell Biochem	WB
Xiaolei Zhang	36382280	Chem Sci	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10430-1-AP (CDK5 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using CDK5 antibody (10430-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 10430-1-AP (CDK5 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 10430-1-AP (CDK5 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).