For Research Use Only

GORASP2 Polyclonal antibody

Catalog Number: 10598-1-AP

Featured Product

60 Publications



Basic Information

Catalog Number: GenBank Accession Number: 10598-1-AP BC007770

Size: GeneID (NCBI): 26003

Source: UNIPROT ID: Rabbit Q9H8Y8

Isotype: Full Name:

golgi reassembly stacking protein 2,

Immunogen Catalog Number: 55kDa

AG0950 Calculated MW:

55 kDa Observed MW: 59 kDa

Applications

Tested Applications: IF/ICC, IHC, IP, WB, ELISA

Cited Applications: WB, IF, IHC

Species Specificity: human, mouse, rat

Cited Species:

human, mouse, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: COLO 320 cells, rat testis tissue, HeLa cells, Raji

Purification Method:

WB 1:500-1:3000

protein lysate

IHC 1:20-1:200

IF 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

cells

IP: HeLa cells,

IHC: human colon cancer tissue,

IF: HepG2 cells,

Background Information

GORASP2 (Golgi reassembly-stacking protein 2, or GRS2) has multiple phosphorylation sites and can be phosphorylated by MKK/ERK pathway during mitoisis. GORASP2 plays a role in the assembly and membrane stacking of the Golgi cisternae, and in the process by which Golgi stacks reform after mitotic breakdown.GORASP2 is supposed to regulate the intracellular transport and presentation of a defined set of transmembrane proteins, such as transmembrane TGFA.

Notable Publications

Author	Pubmed ID	Journal	Application
Jennifer Y Liu	34533192	Biol Open	IF
Xihua Yue	34493279	BMC Biol	WB
Myun Hwa Dunlop	28874656	Nat Commun	

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10598-1-AP (GORASP2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




WB result of GORASP2 antibody (10598-1-AP; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-GORASP2 transfected Hela cells.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using GORASP2 antibody (10598-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL555-Phalloidin (orange).



Immunohistochemical analysis of paraffinembedded human colon cancer using 10598-1-AP (GORASP2 antibody) at dilution of 1:100 (under 10x lens).



IP result of anti-GORASP2 (IP:10598-1-AP, 3ug; Detection:10598-1-AP 1:700) with HeLa cells lysate 1320ug.