## For Research Use Only

# RHOA Polyclonal antibody

Catalog Number: 10749-1-AP

**Featured Product** 

**127 Publications** 

BC005976

GeneID (NCBI):

**UNIPROT ID:** 

P61586 Full Name:



**Basic Information** 

Catalog Number: 10749-1-AP

Isotype:

Size:  $600 \mu g/ml$ Source: Rabbit

Immunogen Catalog Number:

AG1141

Observed MW:

ras homolog gene family, member A Calculated MW: 22 kDa

GenBank Accession Number:

22 kDa

**Applications** 

**Tested Applications:** IF/ICC, WB, ELISA Cited Applications: IF, IHC, WB Species Specificity: human, mouse, rat

**Cited Species:** human, rat, mouse, pig Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000 IF 1:200-1:800

**Purification Method:** 

Positive Controls:

WB: HL-60 cells, HeLa cells, MCF-7 cells, PC-12 cells, rat brain tissue, mouse brain tissue, HUVEC cells, NIH/3T3 cells, fetal human brain tissue

IF: HeLa cells,

# **Background Information**

RhoA is a member of the Rho family of small GTPases, which cycle between inactive GDP-bound and active GTPbound states and function as molecular switches in signal transduction cascades. Rho proteins promote reorganization of the actin cytoskeleton and regulate cell shape, attachment, and motility. Overexpression of RhoA is associated with tumor cell proliferation and metastasis. RhoA signalling is critical to many cellular processes including migration, mechanotransduction, and is often disrupted in carcinogenesis.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Lichan Mao	36247873	Anal Cell Pathol (Amst)	WB
Kai Cui	27678468	Asian J Androl	WB,IHC
Jing Wang	26409448	Tumour Biol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 10749-1-AP (RHOA antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 10749-1-AP (RHOA antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using RHOA antibody (10749-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).