For Research Use Only

PRG2 Polyclonal antibody

Catalog Number: 10766-1-AP

2 Publications



Purification Method:

IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: 10766-1-AP GenBank Accession Number: BC005929 GeneID (NCBI): 5553 UNIPROT ID:

UNIPROT ID: P13727 Full Name:

Immunogen Catalog Number:

AG1150

Size: 400 µg/ml

Source: Rabbit

Isotype:

proteoglycan 2, bone marrow (natural killer cell activator, eosinophil granule major basic protein)

Calculated MW: 25 kDa

Applications

Tested Applications:

IHC,ELISA

Cited Applications:

IF, IHC

Species Specificity:

human Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

IHC : human breast cancer tissue,

Positive Controls:

Background Information

PRG2, also known as BMPG, has 222 amino acid residues with a 16 N-terminal signal sequence, so the mature form has 206 amino acid residues. High levels of PRG2 are present in placenta and pregnancy serum, where it exists as a complex with several other proteins including pregnancy-associated plasma protein A (PAPPA), angiotensinogen (AGT), and C3dg. This protein can be cleaved into eosinophil granule major basic protein (MBP), which is the major constituent of the crystalline core of the eosinophil granule. MBP is highly basic and toxic to both mammalian cells and parasites, and it functions in the defense against invading parasites (PMID: 21270431).

Notable Publications

Author	Pubmed ID	Journal	Application
Jonathan G Medernach	37497566	Allergy	IF
Ohtomo Kaori K	20826153	Exp Eye Res	IHC

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10766-1-AP (PRG2 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10766-1-AP (PRG2 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).