For Research Use Only

ACOT11 Polyclonal antibody

Catalog Number: 10776-1-AP



Purification Method:

WB 1:500-1:2000 IHC 1:20-1:200

IF 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 10776-1-AP BC001517

Size: GeneID (NCBI): 26027

Source: UNIPROT ID: Rabbit Q8WXI4

Isotype: Full Name:

IgG acyl-CoA thioesterase 11

Immunogen Catalog Number: Calculated MW:
AG1176 68.4 kDa
Observed MW:

68-70 kDa

Applications

Tested Applications: WB,IF/ICC,IHC,ELISA Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: DU 145 cells, PC-3 cells

IHC: human prostate cancer tissue,

IF: A431 cells, U2OS cells

Background Information

ACOT11(Acyl-coenzyme A thioesterase 11) is also named as BFIT, KIAAO707, THEA.BFIT is mapped to syntenic regions of chromosomes 1 (human) and 4 (mouse) associated with body fatness and diet-induced obesity, potentially linking a deficit of BFIT activity with exacerbation of these traits and supports the transition of this tissue towards increased metabolic activity, probably through alteration of intracellular fatty acyl-CoA concentration(PMID:11696000).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed A431 cells using 10776-1-AP (ACOT11 antibody), at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human prostate cancer using 10776-1-AP (ACOT11 antibody) at dilution of 1:100 (under 10x lens).



Various lysates were subjected to SDS PAGE followed by western blot with 10776-1-AP (ACOT11 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.