

# ATF4 Polyclonal antibody

Catalog Number: 10835-1-AP

Featured Product

357 Publications

## Basic Information

## Catalog Number:

10835-1-AP

## Size:

1000 µg/ml

## Source:

Rabbit

## Isotype:

IgG

## Immunogen Catalog Number:

AG1279

## GenBank Accession Number:

BC022088

## GeneID (NCBI):

468

## UNIPROT ID:

P18848

## Full Name:

activating transcription factor 4 (tax-responsive enhancer element B67)

## Calculated MW:

39 kDa

## Observed MW:

45-50 kDa

## Purification Method:

Antigen affinity purification

## Recommended Dilutions:

WB 1:500-1:1000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF 1:200-1:800

## Applications

## Tested Applications:

FC, IF/ICC, IHC, IP, WB, ELISA

## Cited Applications:

ChIP, CoIP, FC, IF, IHC, IP, WB

## Species Specificity:

human, rat, mouse

## Cited Species:

human, rat, zebrafish, hamster, pig

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Positive Controls:

WB : A431 cells, HEK-293 cells, rat brain tissue

IP : HEK-293 cells,

IHC : human breast cancer tissue, mouse colon tissue

IF : Tunicamycin treated HeLa cells, HepG2 cells, HeLa cells

## Background Information

ATF4 is a transcription factor, that accumulates predominantly in osteoblasts, where it regulates terminal osteoblast differentiation and bone formation [PMID: 19016586]. As a basic leucine-zipper (bZip) transcription factor, ATF4 can regulate amino acid metabolism, cellular redox state, and anti-stress responses. It also regulates age-related and diet-induced obesity and glucose homeostasis in mammals, and has conserved metabolic functions in flies [PMID: 19726872]. Due to its location at chromosome 22q13, a region linked to schizophrenia, ATF4 is considered as a positional candidate gene for schizophrenia [PMID: 18163433]. Otherwise, since ATF4 is induced by tumour microenvironmental factors, and regulates processes relevant to cancer progression, it might serve as a potential therapeutic target in cancer. Endogenous ATF4 protein has a molecular mass of 50 kDa [PMID: 17726049]. This antibody is a rabbit polyclonal antibody raised against full length human ATF4 antigen. The antibody recognizes the 45-50 kDa ATF4 protein and its phosphorylated forms (50 kDa). ATF4 can bind DNA as a homodimer and as a heterodimer. ATF4 is ubiquitinated by SCF (BTRC) in response to mTORC1 signal, followed by proteasomal degradation and leading to down-regulate expression of SIRT4, so the molecular weight of ATF4 may be 70 kDa.

## Notable Publications

Author	Pubmed ID	Journal	Application
Theresia Sarabhai	36178534	Diabetologia	WB
Junxia Hu	31580970	Biomed Pharmacother	WB
Jing Sun	34650437	Front Pharmacol	WB

## Storage

## Storage:


Store at -20°C. Stable for one year after shipment.

## Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10835-1-AP (ATF4 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 10835-1-AP (ATF4 antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




A431 cells were subjected to SDS PAGE followed by western blot with 10835-1-AP (ATF4 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




IP result of anti-ATF4 (IP:10835-1-AP, 3ug; Detection:10835-1-AP 1:400) with HEK-293 cells lysate 6000ug.



Immunohistochemical analysis of paraffin-embedded mouse colon tissue slide using 10835-1-AP (ATF4 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Tunicamycin treated HeLa cells using ATF4 antibody (10835-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



1X10<sup>6</sup> HepG2 cells were intracellularly stained with 0.2 ug Anti-Human ATF4 (10835-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).