For Research Use Only

DACH1 Polyclonal antibody

Catalog Number:10914-1-AP

Featured Product

69 Publications




Basic Information	Catalog Number: 10914-1-AP	GenBank Accession Number: BC021219	Purification Method: Antigen affinity purification	
	Size:	GeneID (NCBI):	Recommended Dilutions:	
	600 µg/ml	1602	WB 1:5000-1:50000	
	Source: Rabbit	UNIPROT ID: Q9UI36	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
	Isotype: IgG	Full Name: dachshund homolog 1 (Drosophila)	IHC 1:50-1:500 IF 1:200-1:800	
				Immunogen Catalog Number: AG1354
		Observed MW: 97-110 kDa		
	Applications	Tested Applications:	Positive Cor	Positive Controls:
WB, IP, IF/ICC, IHC, ELISA Cited Applications:		WB: H9C2 ce	WB : H9C2 cells, HEK-293 cells	
WB, IP, IF, IHC, ChIP		IP : HEK-293	IP : HEK-293 cells,	
Species Specificity: human, mouse, rat		IHC : human tissue	IHC : human oesophagus cancer tissue, mouse brain tissue	
Cited Species: IF : HEK-293 cells,				
human, rat, mouse, zebrafish, pig				
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
	DACH1, a homologue of the Drosophila dachshund gene, is a key regulator of cell fate determination during eye, leg, and brain development in the fly. Through interacting with NCoR and Smad4, DACH1 is able to inhibit the transforming growth factor-beta (TGF-beta) signaling pathway. DACH1 can inhibit breast cancer cellular proliferation via cyclin D1, suggesting a possible role in tumor suppression. Additionally, DACH1 plays an importa role in negative regulation of RANKL (Receptor activator of NF-kappaB ligand) gene expression in marrow stromal/preosteoblast cells, and loss of DACH1 expression might be involved in endometrial cancer progression. Four isoforms of DACH1 are produced by alternative splicing. This antibody is a rabbit polyclonal antibody raised against residues near the C terminus of human DACH1.			
Background Information	transforming growth factor-beta (proliferation via cyclin D1, sugger role in negative regulation of RAN stromal/preosteoblast cells, and l Four isoforms of DACH1 are produ	sting a possible role in tumor suppressio IKL (Receptor activator of NF-kappaB ligg oss of DACH1 expression might be invol ced by alternative splicing. This antiboo	n inhibit breast cancer cellular n. Additionally, DACH1 plays an importa and) gene expression in marrow ved in endometrial cancer progression.	
	transforming growth factor-beta (proliferation via cyclin D1, sugges role in negative regulation of RAN stromal/preosteoblast cells, and l Four isoforms of DACH1 are produ against residues near the C termin	sting a possible role in tumor suppressio IKL (Receptor activator of NF-kappaB ligg oss of DACH1 expression might be invol ced by alternative splicing. This antiboo	n inhibit breast cancer cellular n. Additionally, DACH1 plays an importa and) gene expression in marrow ved in endometrial cancer progression.	
	transforming growth factor-beta (proliferation via cyclin D1, sugger role in negative regulation of RAN stromal/preosteoblast cells, and I Four isoforms of DACH1 are produ against residues near the C termin	sting a possible role in tumor suppressio IKL (Receptor activator of NF-kappaB lig: oss of DACH1 expression might be invol iced by alternative splicing. This antiboo nus of human DACH1.	n inhibit breast cancer cellular n. Additionally, DACH1 plays an import and) gene expression in marrow ved in endometrial cancer progression. ly is a rabbit polyclonal antibody raised	
	transforming growth factor-beta (proliferation via cyclin D1, sugger role in negative regulation of RAN stromal/preosteoblast cells, and l Four isoforms of DACH1 are produ against residues near the C termin Author Fernando Bonet	sting a possible role in tumor suppressio IKL (Receptor activator of NF-kappaB ligs oss of DACH1 expression might be invol iced by alternative splicing. This antiboo hus of human DACH1. Pubmed ID Journal	n inhibit breast cancer cellular n. Additionally, DACH1 plays an importa and) gene expression in marrow ved in endometrial cancer progression. ly is a rabbit polyclonal antibody raised Application IF	
	transforming growth factor-beta (proliferation via cyclin D1, sugger role in negative regulation of RAN stromal/preosteoblast cells, and I Four isoforms of DACH1 are produ against residues near the C termin Author Fernando Bonet Masahiro Okabe	sting a possible role in tumor suppressio IKL (Receptor activator of NF-kappaB ligs oss of DACH1 expression might be invol iced by alternative splicing. This antiboo hus of human DACH1. Pubmed ID Journal 30204931 Dev Dyn	n inhibit breast cancer cellular n. Additionally, DACH1 plays an importa and) gene expression in marrow ved in endometrial cancer progression. ly is a rabbit polyclonal antibody raised Application IF	
	transforming growth factor-beta (proliferation via cyclin D1, sugger role in negative regulation of RAN stromal/preosteoblast cells, and I Four isoforms of DACH1 are produ against residues near the C termin Author Fernando Bonet Masahiro Okabe	sting a possible role in tumor suppressio IKL (Receptor activator of NF-kappaB lig: oss of DACH1 expression might be invol iced by alternative splicing. This antiboo nus of human DACH1. Pubmed ID Journal 30204931 Dev Dyn 30379099 Am J Physiol Renal	n inhibit breast cancer cellular n. Additionally, DACH1 plays an import and) gene expression in marrow ved in endometrial cancer progression. ly is a rabbit polyclonal antibody raised Application IF	
Background Information Notable Publications Storage	transforming growth factor-beta (proliferation via cyclin D1, sugger role in negative regulation of RAN stromal/preosteoblast cells, and I Four isoforms of DACH1 are produ against residues near the C termin Author Fernando Bonet Masahiro Okabe	sting a possible role in tumor suppressio IKL (Receptor activator of NF-kappaB lig: oss of DACH1 expression might be invol iced by alternative splicing. This antiboo nus of human DACH1. Pubmed ID Journal 30204931 Dev Dyn 30379099 Am J Physiol Renal 24149323 Epigenetics	n inhibit breast cancer cellular n. Additionally, DACH1 plays an importa and) gene expression in marrow ved in endometrial cancer progression. ly is a rabbit polyclonal antibody raised Application IF	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10914-1-AP (DACH1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




WB result of DACH1 (10914-1-AP, 1:4000) with sicontrol and si-DACH1 transfected HEK293 cells.



IP result of anti-DACH1 (IP:10914-1-AP, 4ug; Detection:10914-1-AP 1:1000) with HEK-293 cells lysate 6000ug.



Immunohistochemical analysis of paraffinembedded human oesophagus cancer tissue slide using 10914-1-AP (DACH1 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human oesophagus cancer tissue slide using 10914-1-AP (DACH1 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using DACH1 antibody (10914-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).