## For Research Use Only

## Nurr1/NR4A2 Polyclonal antibody

Catalog Number: 10975-2-AP 17 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 10975-2-AP BC009288 GeneID (NCBI): Size: 700 μg/ml 4929 **UNIPROT ID:** Source: Rabbit P43354 Full Name: Isotype:

nuclear receptor subfamily 4, group A,

member 2 Immunogen Catalog Number: AG1418 Calculated MW: 66 kDa

> Observed MW: 66 kDa

**Applications** 

**Tested Applications:** WB, IHC, ELISA **Cited Applications:** WB, IF, IHC, ChIP Species Specificity: human

**Cited Species:** human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: COLO 320 cells, fetal human brain tissue, SH-

**Purification Method:** 

WB 1:1000-1:4000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

SY5Y cells, A431 cells, HUVEC cells

IHC: human gliomas tissue,

**Background Information** 

NR4A2 belongs to the orphan nuclear receptor (NR) family referred to as NR4A family, which have been implicated in cell cycle regulation, apoptosis, inflammation, metabolism and more recently in carcinogenesis [PMID:22583411]. NR4A2 also has a role in the expression of several proteins that are necessary for the synthesis and regulation of DA, such as tyrosine hidroxilase, DA transporter, vesicular monoamine transporter 2, and cRET [PMID:22294735]. Act as a transcriptional regulator, NR4A2 is essential for the differentiation and maintenance of meso-diencephalic dopaminergic (mdDA) neurons during development. It is crucial for expression of a set of genes such as SLC6A3, SLC18A2, TH and DRD2 which are essential for development of mdDA neurons [PMID: 17184956].

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Xiaoxuan Zhai	36184541	Int Heart J	WB
Guoqing Sun	36138991	Brain Sci	WB
Jian Huang	36089218	Brain Behav Immun	WB,IF

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 10975-2-AP (Nurr1/NR4A2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 10975-2-AP (Nurr1/NR4A2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.