## For Research Use Only

## CDK4 Polyclonal antibody

Catalog Number: 11026-1-AP

Featured Product

**340 Publications** 



**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 11026-1-AP
 BC010153

 Size:
 GeneID (NCBI):

 450 µg/ml
 1019

 Source:
 UNIPROT ID:

 Rabbit
 P11802

Rabbit P11802 Isotype: Full Name:

IgG cyclin-dependent kinase 4

Immunogen Catalog Number: Calculated MW:

AG1505 34 kDa

Observed MW: 30-34 kDa **Purification Method:** 

Antigen affinity purification

Recommended Dilutions: WB 1:1000-1:8000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF 1:50-1:500

**Applications** 

**Tested Applications:** 

FC, IF/ICC, IHC, IP, WB, ELISA

Cited Applications: CoIP, IF, IHC, IP, RIP, WB Species Specificity: human, rat, mouse Cited Species:

human, rat, sheep, mouse, zebrafish, pig, canine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Positive Controls:** 

WB: HeLa cells, rat pancreas tissue, HepG2 cells, MCF-7 cells, NIH/3T3 cells

IP: HeLa cells,

IHC: human lung cancer tissue, human breast cancer

tissue

IF: MCF-7 cells,

**Background Information** 

Cyclin-dependent kinase-4 (CDK4) is a protein-serine kinase involved in the cell cycle. It is essential for the G1- to S-phase transition during the cell cycle and its expression is primarily controlled at the transcriptional level(PMID:17253961). CCND1-CDK4 axis is not only critical in glial tumor cells but also in stromal-derived cells in the surrounding tumor microenvironment that are vital to sustain tumor outgrowth(PMID:21844184).

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Yuanyuan Bai	36213833	J Oncol	WB
Taiwei Wang	36169181	Oncol Rep	WB
Yue Wang	27690088	Nutrients	WB

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11026-1-AP (CDK4 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 11026-1-AP (CDK4 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 11026-1-AP (CDK4 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




WB result of CDK4 antibody (11026-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CDK4 transfected HeLa cells.



IP result of anti-CDK4 (IP:11026-1-AP, 3ug; Detection:11026-1-AP 1:1000) with HeLa cells lysate 4000ug.



1X10^6 MCF-7 cells were stained with 0.2ug CDK4 antibody (11026-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using CDK4 antibody (11026-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).