

c-SRC Polyclonal antibody

Catalog Number: 11097-1-AP

66 Publications

Basic Information

Catalog Number:

11097-1-AP

Size:

800 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1575

GenBank Accession Number:

BC011566

GeneID (NCBI):

6714

ENSEMBL Gene ID:

ENSG00000197122

UNIPROT ID:

P12931

Full Name:

v-src sarcoma (Schmidt-Ruppin A-2)
viral oncogene homolog (avian)

Calculated MW:

60 kDa

Observed MW:

55 kDa, 60 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 µg for 1.0-3.0 mg of total
protein lysate

IHC 1:100-1:400

IF 1:10-1:100

Applications

Tested Applications:

FC, IF/ICC, IHC, IP, WB, ELISA

Cited Applications:

WB, IP, IF, IHC

Species Specificity:

human, mouse, rat, hamster

Cited Species:

human, rat, mouse

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : HUVEC cells, mouse brain tissue, COLO 320 cells,
Jurkat cells, A431 cells, CHO cells, SH-SY5Y cells

IP : SH-SY5Y cells,

IHC : human breast cancer tissue, human brain tissue,
human spleen tissue

IF : A431 cells,

Background Information

SRC, also named as SRC1 and p60-Src, belongs to the protein kinase superfamily, Tyr protein kinase family and SRC subfamily. It is a non-receptor protein tyrosine kinase that plays pivotal roles in numerous cellular processes such as proliferation, migration, and transformation. In concert with PTK2B, SRC plays an important role in osteoclastic bone resorption. It promotes energy production in osteoclasts by activating mitochondrial cytochrome C oxidase. SRC can be detected as 60 kDa and 55 kDa (PMID: 15047935, 21461171).

Notable Publications

Author	Pubmed ID	Journal	Application
YanHua Fan	36174847	Fitoterapia	WB
Lianhong Yin	29033837	Front Pharmacol	WB
Jiao Wu	34517345	Aging (Albany NY)	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

**This product is exclusively available under Proteintech
Group brand and is not available to purchase from any
other manufacturer.**


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11097-1-AP (SRC antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




IP result of anti-c-SRC (IP:11097-1-AP, 5ug; Detection:11097-1-AP 1:1000) with SH-SY5Y cells lysate 2900ug.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 11097-1-AP (c-SRC Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 11097-1-AP (c-SRC Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of A431 cells using 11097-1-AP (SRC antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10⁶ SH-SY5Y cells were stained with 0.2ug c-SRC antibody (11097-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.



Jurkat cells were subjected to SDS PAGE followed by western blot with 11097-1-AP (SRC antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.