For Research Use Only

Aconitase 2 Polyclonal antibody

Catalog Number:11134-1-AP

15 Publications



Basic Information

Catalog Number: 11134-1-AP Size:

 $600 \mu g/ml$

Source: Rabbit Isotype:

aconitase 2, mitochondrial

Calculated MW: Immunogen Catalog Number: AG1590 85 kDa

Observed MW:

85 kDa

BC014092

GeneID (NCBI):

UNIPROT ID:

Q99798 Full Name:

GenBank Accession Number:

Purification Method:

Antigen affinity purification Recommended Dilutions: WB 1:2000-1:12000

IHC 1:20-1:200 IF 1:50-1:500

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA Cited Applications:

IF. WB

Species Specificity: human, mouse, rat **Cited Species:** human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

WB: mouse heart tissue, human heart tissue, human liver tissue, human skeletal muscle tissue, HeLa cells, mouse skeletal muscle tissue, rat heart tissue, rat skeletal muscle tissue

IHC: human lung cancer tissue, human prostate cancer tissue

IF: HeLa cells.

Background Information

ACO2(aconitate hydratase, mitochondrial) is also named as citrate hydro-lyase and belongs to the aconitase/IPM isomerase family. It plays a key function in cellular energy production, and loss of its activity has a major impact on cellular and organismal survival. Western blot shows two bands of 83 kDa and 40 kDa. The 40 kDa fragment decreases with age and oxidative stress, whereas other fragmentation products with molecular weights between 40 and 83 kDa increased with age and MnSOD(mitochondrial manganese superoxide dismutase) deficiency(PMID:12459471). Defects in ACO2 are the cause of infantile cerebellar-retinal degeneration (ICRD).

Notable Publications

Author	Pubmed ID	Journal	Application
Minghao Yuan	36186135	Aging Dis	WB
Chiung-Mei Chen	29160844	Int J Mol Sci	WB
Bruna Rodrigues Muys	31101765	Cancer Res	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 11134-1-AP (Aconitase 2 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung cancer using 11134-1-AP (Aconitase 2 antibody) at dilution of 1:50 (under 10x lens).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using 11134-1-AP (Aconitase 2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).