

For Research Use Only

# OCT4/POU5F1 Polyclonal antibody

Catalog Number: 11263-1-AP

Featured Product

212 Publications



## Basic Information

### Catalog Number:

11263-1-AP

### Size:

750 µg/ml

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG1794

### GenBank Accession Number:

BC020712

### GeneID (NCBI):

5460

### UNIPROT ID:

Q01860

### Full Name:

POU class 5 homeobox 1

### Calculated MW:

39 kDa

### Observed MW:

50-60 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:1000-1:5000

IHC 1:50-1:500

IF 1:50-1:500

## Applications

### Tested Applications:

FC, IF-P, IHC, WB, ELISA

### Cited Applications:

FC, IP, WB

### Species Specificity:

human, mouse, rat

### Cited Species:

human, goat, rat, mouse, pig

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB** : HEK-293 cells, MCF-7 cells, NCCIT cells, MDA-MB-231 cells, mouse brain tissue, U-251 cells, rat brain tissue, mouse embryo tissue

**IHC** : human renal cell carcinoma tissue, human lung cancer tissue

**IF** : mouse eye tissue,

## Background Information

Octamer-binding protein 4 (OCT4), also known as POU5F1 or OCT3/4, is a member of the POU gene family and encoded by the human gene POU5F1 (POU domain class 5 transcription factor 1). OCT4 is a member of the Octamer class of transcription factors that recognize the 8-base pair consensus motif 5'-ATGCAAAT-3'. Expression of OCT4 is associated with an undifferentiated, pluripotent stem cell phenotype and tumors. OCT4 is also expressed in the developing brain, with the highest levels found in the cortex, olfactory bulb, and the hippocampus.

## Notable Publications

Author	Pubmed ID	Journal	Application
Yang Liu	27685621	Cell Death Dis	WB
Chenlong Li	31558707	Cell Death Dis	WB
XIAOYUE FENG	34528694	Oncol Rep	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11263-1-AP (OCT4/POU5F1 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human renal cell carcinoma tissue slide using 11263-1-AP (OCT4/POU5F1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse eye tissue using OCT4/POU5F1 antibody (11263-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10<sup>6</sup> HEK-293T cells were intracellularly stained with 0.4 ug Anti-Human OCT4/POU5F1 (11263-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).