For Research Use Only

PRMT1 Polyclonal antibody

Catalog Number: 11279-1-AP

Featured Product

8 Publications



Basic Information

Catalog Number: GenBank Accession Number: 11279-1-AP BC019268

Size: GeneID (NCBI): 3276

Source: UNIPROT ID: Rabbit Q99873

Isotype: Full Name:

protein arginine methyltransferase 1

Immunogen Catalog Number: Calculated MW: AG1790 42 kDa

Observed MW: 40-42 kDa

Applications

Tested Applications: WB,IP,IHC,IF/ICC,FC,ELISA

Cited Applications: WB,IP,IHC,IF Species Specificity: human, mouse, rat

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A549 cells, mouse brain tissue, MCF-7 cells, HeLa cells, NIH/3T3 cells, rat brain tissue, Jurkat cells

Purification Method:

WB 1:2000-1:16000

protein lysate

IHC 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

IP: A549 cells,

IHC: human colon tissue,

IF: A431 cells,

Background Information

PRMT1, the major protein arginine methyltransferase in mammals, catalyzes monomethylation and asymmetric dimethylation of arginine side chains in proteins. Human PRMT1 is encoded by the PRMT1 gene located on chromosome 19 (19q13.3) and composed of 12 exons and 11 introns. At the protein level, human PRMT1 shares a high degree of homology with the different members of the PRMT family that is conserved in eukaryotes. Phylogenetic studies based on the methyltransferase domain highlighted that PRMT1 is closely related to PRMT8.

Notable Publications

Author	Pubmed ID	Journal	Application
Zhongwei Li	34775498	Cell Death Dis	WB,IHC
Xin-Ke Yin	33420374	Oncogene	IHC
Yizhen Tian	34919946	Toxicol Appl Pharmacol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




WB result of PRMT1 antibody (11279-1-AP; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PRMT1 transfected A549 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 11279-1-AP (PRMT1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 11279-1-AP (PRMT1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 11279-1-AP (PRMT1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




IP result of anti-PRMT1 (IP:11279-1-AP, 4ug; Detection:11279-1-AP 1:2000) with A549 cells lysate 920 ug.




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 11279-1-AP (PRMT1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 11279-1-AP (PRMT1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using PRMT1 antibody (11279-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 Raji cells were stained with 0.2ug PRMT1 antibody (11279-1-AP, red) and control antibody (blue). Cells were fixed with 4% PFA and permeabilized with 0.1% Triton X-100. Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit $\lg G(H+L)$ with dilution 1:1500.