For Research Use Only

ZBTB7B Polyclonal antibody

Catalog Number: 11341-1-AP

3 Publications



Basic Information

Catalog Number: 11341-1-AP

Size: 800 μ g/ml Source: Rabbit

Isotype: IgG

Immunogen Catalog Number: AG1888

GenBank Accession Number:

BC012070 GeneID (NCBI): 51043 Full Name:

zinc finger and BTB domain containing 7B

Calculated MW: 539 aa, 58 kDa Observed MW: 60-65 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:4000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IF 1:10-1:100

Applications

Tested Applications: IF/ICC, IP, WB, ELISA

Cited Applications: WB, IP, IHC

Species Specificity: human, mouse, rat Cited Species: human, hamster, bovine Positive Controls:

WB: HeLa cells, HepG2 cells

IP: HeLa cells,

IF: Hela cells, MCF-7 cells

Background Information

ZBTB7B belongs to a large family of transcription factors, generally acting as repressors, characterized by a carboxyterminal DNA binding domain made of multiple zinc fingers (four in Thpok) and an amino-terminal BTB-POZ domain that mediates homo- (and possibly hetero-) dimerization [PMID: 17084908]. Zbtb7b is up-regulated by MHC-II-restricted thymocytes during their CD4 differentiation and is a major determinant of CD4 lineage choice. Two properties of ZBTB7B deserve attention. First, although Thpok is expressed in a wide variety of cells, its expression in the thymus is highly lineage-specific: CD4 SP thymocytes (and all CD4 T cells) express Thpok, whereas DP and CD8 SP thymocytes do notSecond, both loss- and gain-of-function experiments indicate that Thpok affects lineage choice but not positive selection [PMID: 15729333, 10550051].

Notable Publications

Author	Pubmed ID	Journal	Application
Jian Sun	34700042	Acta Biomater	WB
Hao-Ming Xu	35761286	J Transl Med	WB,IHC
Jiaxin Wang	38154091	J Agric Food Chem	WB,IP

Storage


Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 11341-1-AP (ZBTB7B antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of Hela cells, using ZBTB7B antibody 11341-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP result of anti-ZBTB7B (IP:11341-1-AP, 4ug; Detection:11341-1-AP 1:3000) with HeLa cells lysate 1320 ug.