For Research Use Only

## ARF4 Polyclonal antibody

Catalog Number:11673-1-AP

Featured Product






| Basic Information      | Catalog Number:<br>11673-1-AP   | GenBank Accession Number:<br>BC003364   | Purification Method:<br>Antigen affinity purification  |
|------------------------|---|---|--|
|                        | Size:   | GenelD (NCBI):  | Recommended Dilutions:   |
|                        | 300 µg/ml   | 378   | WB 1:1000-1:6000   |
|                        | Source:   | UNIPROT ID:   | IP 0.5-4.0 ug for 1.0-3.0 mg of total  |
|                        | Rabbit  | P18085  | protein lysate   |
|                        | Isotype:  | Full Name:  | IHC 1:50-1:500<br>IF 1:50-1:500  |
|                        | lgG   | ADP-ribosylation factor 4   | 11 1.50-1.500  |
|                        | Immunogen Catalog Number:<br>AG2272   | Calculated MW:<br>180 aa, 21 kDa  |  |
|                        |   | Observed MW:<br>16 kDa, 21 kDa  |  |
| Applications           | Tested Applications:<br>FC, IF/ICC, IHC, IP, WB, ELISA  | Positive Controls:  |  |
|                        | Cited Applications:   |   | IT-29 cells, HeLa cells, HepG2 cells, K-562 cells<br>e brain tissue, rat brain tissue  |
|                        | IF, IHC, WB   | IP : He   | pG2 cells, mouse brain tissue  |
|                        | Species Specificity:<br>human, mouse, rat   | IHC :   | numan breast cancer tissue,  |
|                        | Cited Species:  | IF : He   | epG2 cells,  |
|                        | human, mouse  |   |  |
|                        | Note-IHC: suggested antigen retrieval with<br>TE buffer pH 9.0; (*) Alternatively, antigen<br>retrieval may be performed with citrate<br>buffer pH 6.0  |   |  |
|                        |   |   |  |
| Background Information | are members of the ARF family of<br>and regulate GTP/GDP cycle by al<br>ARF family proteins are essential<br>been identified in mammalian ce  | GTP-binding proteins of the Ras s<br>ternating between the active GTP<br>and ubiquitous in eukaryotes. Siv<br>Ils. They function in vesicular traf<br>I ARFs. Their function are not full   | uperfamily, with 20 kDa protein size. ARFs bind<br>-bound and inactive GDP-bound conformations<br>highly conserved members of the family have  |
|                        | are members of the ARF family of<br>and regulate GTP/GDP cycle by al<br>ARF family proteins are essential<br>been identified in mammalian ce<br>in cells. ARF 4 and ARF 5 are class I<br>This antibody can bind ARFs for th   | GTP-binding proteins of the Ras s<br>ternating between the active GTP<br>and ubiquitous in eukaryotes. Siv<br>Ils. They function in vesicular traf<br>I ARFs. Their function are not full   | y and Arf family. ADP-ribosylation factors (ARF<br>uperfamily, with 20 kDa protein size. ARFs bind<br>-bound and inactive GDP-bound conformations<br>highly conserved members of the family have<br>fic and actin remodelling and other bioprocess<br>y described. (PMID: 7759471, PMID: 16042562).<br>Application |
|                        | are members of the ARF family of<br>and regulate GTP/GDP cycle by al<br>ARF family proteins are essential<br>been identified in mammalian ce<br>in cells. ARF4 and ARF5 are class<br>This antibody can bind ARFs for th<br>Author                                     | GTP-binding proteins of the Ras s<br>ternating between the active GTP<br>and ubiquitous in eukaryotes. Siv<br>Ils. They function in vesicular traf<br>I ARFs. Their function are not fully<br>le close sequences.   | uperfamily, with 20 kDa protein size. ARFs bind<br>-bound and inactive GDP-bound conformations<br>highly conserved members of the family have<br>fic and actin remodelling and other bioprocess<br>y described. (PMID: 7759471, PMID: 16042562).   |
| Background Information | are members of the ARF family of<br>and regulate GTP/GDP cycle by al<br>ARF family proteins are essential<br>been identified in mammalian ce<br>in cells. ARF 4 and ARF 5 are class<br>This antibody can bind ARFs for th<br>Author                                   | GTP-binding proteins of the Ras s<br>ternating between the active GTP<br>and ubiquitous in eukaryotes. Siv<br>Ils. They function in vesicular traf<br>I ARFs. Their function are not full<br>te close sequences.<br>Pubmed ID Journal<br>23050017 PLoS One  | uperfamily, with 20 kDa protein size. ARFs bine-<br>bound and inactive GDP-bound conformations<br>highly conserved members of the family have<br>fic and actin remodelling and other bioprocess<br>y described. (PMID: 7759471, PMID: 16042562).<br>Application<br>WB  |
|                        | are members of the ARF family of<br>and regulate GTP/GDP cycle by al<br>ARF family proteins are essential<br>been identified in mammalian ce<br>in cells. ARF4 and ARF5 are class I<br>This antibody can bind ARF5 for th<br>Author I<br>Jain Sachi S<br>Arisa Ochiai | GTP-binding proteins of the Ras s<br>ternating between the active GTP<br>and ubiquitous in eukaryotes. Six<br>ILs. They function in vesicular traf<br>II ARFs. Their function are not full<br>the close sequences.<br>Pubmed ID Journal   | uperfamily, with 20 kDa protein size. ARFs bir<br>-bound and inactive GDP-bound conformation<br>highly conserved members of the family have<br>fic and actin remodelling and other bioprocess<br>y described. (PMID: 7759471, PMID: 16042562)<br>Application<br>WB   |
|                        | are members of the ARF family of<br>and regulate GTP/GDP cycle by al<br>ARF family proteins are essential<br>been identified in mammalian ce<br>in cells. ARF4 and ARF5 are class I<br>This antibody can bind ARF5 for th<br>Author I<br>Jain Sachi S<br>Arisa Ochiai | GTP-binding proteins of the Ras s<br>ternating between the active GTP<br>and ubiquitous in eukaryotes. Six<br>Ils. They function in vesicular traf<br>Il ARFs. Their function are not fully<br>the close sequences.<br>Pubmed ID Journal<br>23050017 PLoS One<br>34523057 Neurochem R<br>26330566 J Biochem | uperfamily, with 20 kDa protein size. ARFs bin<br>-bound and inactive GDP-bound conformations<br>-highly conserved members of the family have<br>fic and actin remodelling and other bioprocess<br>y described. (PMID: 7759471, PMID: 16042562).<br>Application<br>WB<br>es WB                                     |


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11673-1-AP (ARF4 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




WB result of ARF4 antibody (11673-1-AP; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ARF4 transfected HeLa cells.




IP result of anti-ARF4 (IP:11673-1-AP, 3ug; Detection:11673-1-AP 1:600) with mouse brain tissue lysate 5200ug.




IP result of anti-ARF4 (IP:11673-1-AP, 4ug; Detection:11673-1-AP 1:3000) with HepG2 cells lysate 1080 ug.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 11673-1-AP (ARF4 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 11673-1-AP (ARF4 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 11673-1-AP (ARF4 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 K-562 cells were intracellularly stained with 0.4 ug Anti-Human ARF4 (11673-1-AP) and Coralite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).