

For Research Use Only

# STAT5A/B Polyclonal antibody

Catalog Number: 12071-1-AP **7 Publications**



## Basic Information

<b>Catalog Number:</b> 12071-1-AP	<b>GenBank Accession Number:</b> BC020868	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 600 µg/ml	<b>GeneID (NCBI):</b> 6777	<b>Recommended Dilutions:</b> WB 1:1000-1:6000
<b>Source:</b> Rabbit	<b>ENSEMBL Gene ID:</b> ENSG00000173757	IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate
<b>Isotype:</b> IgG	<b>UNIPROT ID:</b> P51692	IHC 1:50-1:500
<b>Immunogen Catalog Number:</b> AG2709	<b>Full Name:</b> signal transducer and activator of transcription 5B	IF 1:10-1:100
	<b>Calculated MW:</b> 787 aa, 90 kDa	
	<b>Observed MW:</b> 90 kDa	

## Applications

<b>Tested Applications:</b> IF/ICC, IHC, IP, WB, ELISA	<b>Positive Controls:</b> WB : HeLa cells, Jurkat cells, K-562 cells, MOLT-4 cells
<b>Cited Applications:</b> IF, IP, WB	IP : HeLa cells,
<b>Species Specificity:</b> human	IHC : human cervical cancer tissue, human lung cancer tissue, human breast cancer tissue
<b>Cited Species:</b> human, rat, mouse	IF : HepG2 cells,
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	

## Background Information

STAT5B belongs to the transcription factor STAT family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. STAT5B carries out a dual function: signal transduction and activation of transcription. STAT5B mediates the signal transduction triggered by various cell ligands, such as IL2, IL4, CSF1, and different growth hormones. It has been shown to be involved in diverse biological processes, such as TCR signaling, apoptosis, adult mammary gland development, and sexual dimorphism of liver gene expression. Although this antibody is a rabbit polyclonal antibody raised against residues near the N terminus of human STAT5B, the homology between this segment and the corresponding segment of human STAT5A is 92%. It shows that this antibody can cross-react with STAT5A.

## Notable Publications

Author	Pubmed ID	Journal	Application
Huan Ge	36306591	Biomed Pharmacother	WB
Furong Zhang	26493931	Toxicol Appl Pharmacol	WB, IP, IF
Dong Ma	27856242	Biochem Biophys Res Commun	WB


## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:  
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12071-1-AP (STAT5A/B antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



IP result of anti-STAT5A/B (IP:12071-1-AP, 4ug; Detection:12071-1-AP 1:4000) with HeLa cells lysate 1720 ug.



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 12071-1-AP (STAT5A/B antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HepG2 cells using 12071-1-AP (STAT5A/B antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).