## For Research Use Only

## SMAD5 Polyclonal antibody

Catalog Number:12167-1-AP

**Featured Product** 

**20 Publications** 



**Basic Information** 

Catalog Number: 12167-1-AP Size:

 $600 \mu g/ml$ Source: Rabbit Isotype:

Immunogen Catalog Number:

AG2809

GenBank Accession Number:

BC009682 GeneID (NCBI): 4090 **UNIPROT ID:** Q99717 Full Name:

SMAD family member 5

Calculated MW: 465 aa, 52 kDa Observed MW: 52 kDa

**Purification Method:** 

Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IF 1:10-1:100

**Applications** 

**Tested Applications:** 

FC, IF/ICC, IP, WB,ELISA Cited Applications: WB. FC

Species Specificity: human, mouse, rat **Cited Species:** 

human, rat, mouse

Positive Controls:

WB: K-562 cells, IP: K-562 cells, IF: MCF-7 cells,

**Background Information** 

SMAD5, also named as MADH5 and JV5-1, is a transcriptional modulator activated by BMP (bone morphogenetic proteins) type 1 receptor kinase. SMAD5 is a receptor-regulated SMAD (R-SMAD). It plays a critical role in the signaling pathway by which TGF-beta inhibits the proliferation of human hematopoietic progenitor cells.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Xueni Wang	32879393	Sci Rep	WB
Jinliang Chen	29715457	Biochem Biophys Res Commun	WB
Wei Peng	35314669	Cell Death Dis	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**




WB result of SMAD5 antibody (12167-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SMAD5 transfected K-562 cells.




K-562 cells were subjected to SDS PAGE followed by western blot with 12167-1-AP (SMAD5 Antibody) at dilution of 1:1000 incubated at 4 degree celsius over night.



Immunofluorescent analysis of MCF-7 cells, using SMAD5 antibody 12167-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP result of anti-SMAD5 (IP:12167-1-AP, 4ug; Detection:12167-1-AP 1:1000) with K-562 cells lysate 3200ug.



1X10^6 MCF-7 cells were stained with 0.2ug SMAD5 antibody (12167-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.